

Practice problems for Linear Algebra Test III.

1. Suppose $L: V \rightarrow W$ is a linear transformation, where the dimension of V is n and the dimension of W is m .

(a) State the rank-nullity theorem for L .

(b) Suppose $n = 7$ and $m = 4$. If L is onto, what is the dimension of the kernel of L ? Explain.

(c) Suppose $V = W$ and L is one-to-one. What else can you say about L ? Explain.

2. Verify the rank-nullity theorem for $L: \mathbf{R}^3 \rightarrow \mathbf{R}^2$, where

$$L \left(\begin{bmatrix} a \\ b \\ c \end{bmatrix} \right) = \begin{bmatrix} a - b + c \\ -a + b - c \end{bmatrix},$$

3. Use determinants and cofactors to find the inverse of the following matrix:

$$\begin{bmatrix} 2 & -1 & 0 \\ 1 & 0 & -2 \\ 3 & 2 & 1 \end{bmatrix}$$

4. Suppose $L: P_2 \rightarrow \mathbf{R}^2$ is defined by

$$L(p(t)) = \begin{bmatrix} p'(1) \\ \int_0^1 p(t) dt \end{bmatrix}.$$

(a) Verify that L is linear.

(b) Let $S = \{t^2 + 1, t + 1, t - 1\}$ be a basis for P_2 , and $T = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ be a basis for \mathbf{R}^2 . Find the matrix representation for L with respect to S and T .

5. Suppose $L: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ is defined by

$$L \left(\begin{bmatrix} a \\ b \end{bmatrix} \right) = \begin{bmatrix} a - 3b \\ 2a + b \end{bmatrix}.$$

(a) Suppose $S = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ is the standard basis for \mathbf{R}^2 . Find the matrix representation for L with respect to S .

(b) Use transition matrices to find the matrix representation for L with respect to the basis $T = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$.

6. Find the determinants of the following matrices:

$$(a) \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \quad (b) \begin{bmatrix} 1 & 2 & -1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix} \quad (c) \begin{bmatrix} 1 & 7 & 4 & -3 \\ 2 & -9 & 0 & 1 \\ 0 & 2 & 0 & 0 \\ 1 & 0 & 1 & 1 \end{bmatrix} \quad (d) \begin{bmatrix} 3 & 5 & 2 & 9 & 0 \\ 5 & 3 & 9 & 5 & 1 \\ 0 & 1 & 4 & 2 & 9 \\ 1 & 8 & 5 & 2 & 0 \\ 5 & 3 & 9 & 5 & 1 \end{bmatrix}$$