

ON THE VANISHING RANGES FOR THE COHOMOLOGY OF FINITE GROUPS OF LIE TYPE

CHRISTOPHER P. BENDEL, DANIEL K. NAKANO, AND CORNELIUS PILLEN

ABSTRACT. Let $G(\mathbb{F}_q)$ be a finite Chevalley group defined over the field of $q = p^r$ elements, and k be an algebraically closed field of characteristic $p > 0$. A fundamental open and elusive problem has been the computation of the cohomology ring $H^\bullet(G(\mathbb{F}_q), k)$. In this paper we determine initial vanishing ranges which improves upon known results. For root systems of type A_n and C_n , the first non-trivial cohomology classes are determined when p is larger than the Coxeter number. In the process we make the use of techniques involving line bundle cohomology for the flag variety G/B and its relation to combinatorial data from Kostant Partition Functions.

1 Introduction

1.1 Let G be a simple algebraic group over an algebraically closed field k of prime characteristic $p > 0$ which is split over the prime field \mathbb{F}_p . Let $\text{Fr} : G \rightarrow G$ denote the Frobenius map and set $q = p^r$. The fixed points of the r th iterate of the Frobenius map, denoted $G(\mathbb{F}_q)$, is a finite Chevalley group. An outstanding open problem of major interest for algebraists and topologists has been to determine the cohomology ring $H^\bullet(G(\mathbb{F}_q), k)$ ¹. In 2005, during a talk at an Oberwolfach conference, Friedlander mentioned that so little is known about this computation that it is not even known in which degree the first non-trivial cohomology class occurs.

Our paper aims to address this fundamental question by investigating two problems:

(1.1.1) Determining Vanishing Ranges: Finding $D > 0$ such that the cohomology groups $H^i(G(\mathbb{F}_q), k) = 0$ for $0 < i < D$.

(1.1.2) Locating the First Non-Trivial Cohomology Class: In many instances in conjunction with the aforementioned problem, we will find a D such that $H^i(G(\mathbb{F}_q), k) = 0$ for $0 < i < D$ and $H^D(G(\mathbb{F}_q), k) \neq 0$. A D satisfying this property will be called a *sharp bound*.

There have been earlier results in the 1970s and 80s addressing (1.1.1). Quillen [Q] showed that $H^i(GL_n(\mathbb{F}_q), k) = 0$ for all $0 < i < r(p - 1)$ and all n . In that work, he noted that the arguments showed for any G as above, there exists a constant C depending on

Date: May 2009.

Research of the first author was supported in part by NSF grant DMS-0400558.

Research of the second author was supported in part by NSF grant DMS-0654169.

2000 *Mathematics Subject Classification.* Primary 20J06; Secondary 20G10.

¹In the cross-characteristic situation (i.e., if $\text{char } k = l$ and $\text{gcd}(l, p) = 1$) much more is known about the cohomology of $H^\bullet(G(\mathbb{F}_q), k)$ (cf. [AM, Chapter VII]). In fact in many cases in the non-describing characteristic the cohomology (including the ring structure) is completely determined.

the root system such that $H^i(G(\mathbb{F}_q), k) = 0$ for $0 < i < r \cdot C$. However, no explicit value of C is given except for $G = SL_2$ (and p odd) in which case one can take $C = (p - 1)/2$. Furthermore, it was not determined whether these vanishing ranges were sharp. Indeed, in the case of SL_2 , one can see from work of Carlson [C] that these bounds are not sharp in general. Quillen's original work arose in the context of certain K -theory computations. Friedlander [F] later used K -theoretic techniques to find vanishing ranges for more general reductive groups. Later work of Hiller [H] extended Friedlander's result and found vanishing ranges for groups of all types. In [FP] Friedlander and Parshall found a sharp bound for the Borel subgroup $B(\mathbb{F}_q)$ of $GL_n(\mathbb{F}_q)$. Since that time, few if any results have been obtained in this direction.

1.2 The strategy in addressing (1.1.1) and (1.1.2) will entail using new and powerful techniques developed by the authors which relate $H^i(G(\mathbb{F}_q), k)$ to extensions over G via a truncated version of the induction functor (cf. [BNP1, BNP2, BNP3, BNP5, BNP6]). An outline of the overall strategy is presented in the diagram below. For the purposes in this paper we will use a non-truncated induction functor $\mathcal{G}_r(-)$. We demonstrate that when applied to the trivial module k , $\mathcal{G}_r(k)$ has a filtration with factors of the form $H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}$ (cf. Proposition 2.4). The G -cohomology of these factors can be analyzed by using the Lyndon-Hochschild-Serre (LHS) spectral sequence involving the first Frobenius kernel G_r (cf. Section 3.1). In particular for $r = 1$, we can apply the results of Kumar-Lauritzen-Thomsen [KLT] to bound the dimension of the cohomology group $H^\bullet(G(\mathbb{F}_p), k)$ from above (cf. Theorem 3.3). The upper bound on the dimension involves the combinatorics of the well-studied Kostant Partition Function. This reduces the question of the vanishing of the finite group cohomology to a question involving the combinatorics of the underlying root system Φ .

$$\begin{array}{c}
 \text{Induction} \\
 \text{Functor} \\
 \\
 H^i(G(\mathbb{F}_q), k) \implies H^i(G, \mathcal{G}_r(k)) \\
 \\
 \Downarrow \text{ Filtrations} \\
 H^i(G, H^0(\lambda) \otimes H(\lambda^*)^{(r)}) \implies H^i(G_1, H^0(\lambda)) \implies \text{Root Combinatorics.} \\
 \\
 \begin{array}{cc}
 \text{LHS Spectral} & \text{Kostant Partition} \\
 \text{Sequences} & \text{Functions}
 \end{array}
 \end{array}$$

More specifically, for a group G of classical type and $H^i(G(\mathbb{F}_q), k)$, under the assumption that $p > h$ (the Coxeter number), in Theorem 4.4, we identify a vanishing range which improves upon (in almost all cases) the ranges of [H]. Furthermore, in type C_n , we identify a *sharp vanishing bound* which addresses (1.1.2) (cf. Theorem 5.4). In type A_n , we identify sharp vanishing bounds for $r = 1$ (cf. Theorem 6.13), and for larger r such sharp vanishing bounds are provided in the generic situation (cf. Theorem 6.14).

The sections of the paper are outlined as follows. In Section 2, we review our previous work and develop the necessary cohomological tools related to induction functors which will be used to determine vanishing ranges. In Section 3, we present some further cohomological properties relating extensions over G with those over the Frobenius kernel G_r . In Section 4, our general vanishing bounds are presented. Finally, Sections 5 and 6 deal with the special cases of root systems of types C_n and A_n respectively. The authors in future work plan to address (1.1.1) and (1.1.2) in the case of the remaining classical groups and the exceptional groups.

2 Relating $G(\mathbb{F}_q)$ and G

2.1 Notation. Throughout this paper, we will follow the basic conventions provided in [J]. Let G be a simple simply connected algebraic group scheme which is defined and split over the finite field \mathbb{F}_p with p elements, and let k be a field of characteristic p . For $r \geq 1$, let $G_r := \ker F^r$ be the r th Frobenius kernel of G and $G(\mathbb{F}_q)$ be the associated finite Chevalley group. Let T be a maximal split torus and Φ be the root system associated to (G, T) . The positive (resp. negative) roots are Φ^+ (resp. Φ^-), and Δ is the set of simple roots. Let B be a Borel subgroup containing T corresponding to the negative roots and U be the unipotent radical of B . For a given root system of rank n , the simple roots will be denoted by $\alpha_1, \alpha_2, \dots, \alpha_n$. We will adhere to the Bourbaki ordering of simple roots. In particular, for type B_n , α_n denotes the unique short simple root and for type C_n , α_n denotes the unique long simple root. The longest (positive) root will be denoted $\tilde{\alpha}$, and for root systems with multiple root lengths, the longest short root will be denoted α_0 . Let W denote the Weyl group associated to Φ , and, for $w \in W$, let $\ell(w)$ denote the length of the word.

Let \mathbb{E} be the Euclidean space associated with Φ , and the inner product on \mathbb{E} will be denoted by $\langle \cdot, \cdot \rangle$. Let $\alpha^\vee = 2\alpha/\langle \alpha, \alpha \rangle$ be the coroot corresponding to $\alpha \in \Phi$. In this case, the fundamental weights (basis dual to $\alpha_1^\vee, \alpha_2^\vee, \dots, \alpha_n^\vee$) will be denoted by $\omega_1, \omega_2, \dots, \omega_n$. Let $X(T)$ be the integral weight lattice spanned by these fundamental weights. The set of dominant integral weights is denoted by $X(T)_+$. For a weight $\lambda \in X(T)$, set $\lambda^* := -w_0\lambda$ where w_0 is the longest word in the Weyl group W . By $w \cdot \lambda := w(\lambda + \rho) - \rho$ we denote the “dot” action of W on $X(T)$, with ρ being the half-sum of the positive roots. For $\alpha \in \Delta$, $s_\alpha \in W$ denotes the reflection in the hyperplane determined by α .

For a G -module M , let $M^{(r)}$ be the module obtained by composing the underlying representation for M with F^r . Moreover, let M^* denote the dual module. For $\lambda \in X(T)_+$, let $H^0(\lambda) := \text{ind}_B^G \lambda$ be the induced module and $V(\lambda) := H^0(\lambda^*)^*$ be the Weyl module of highest weight λ .

2.2 We record two observations on roots that will be used at various points in the paper:

Observation (A). If $\beta \in \Phi^+$ with $\beta \neq \tilde{\alpha}$, then $\langle \beta, \tilde{\alpha}^\vee \rangle \in \{0, 1\}$.

Observation (B). If $w \in W$ admits a reduced expression $w = s_{\beta_1} s_{\beta_2} \dots s_{\beta_m}$ with $\beta_i \in \Delta$ and $m = \ell(w)$, then

$$-w \cdot 0 = \beta_1 + s_{\beta_1}(\beta_2) + s_{\beta_1} s_{\beta_2}(\beta_3) + \dots + s_{\beta_1} s_{\beta_2} \dots s_{\beta_{m-1}}(\beta_m).$$

Moreover, this is the unique way in which $-w \cdot 0$ can be expressed as a sum of distinct positive roots.

2.3 The Induction Functor. Set $\mathcal{G}_r(k) := \text{ind}_{G(\mathbb{F}_q)}^G(k)$. While this G -module is infinite dimensional, it provides a potential way to relate extensions over $G(\mathbb{F}_q)$ with extensions over G . Indeed, by Generalized Frobenius Reciprocity and the fact that $G/G(\mathbb{F}_q)$ is affine, we have the following key observation.

Proposition . *Let M, N be rational G -modules. Then, for all $i \geq 0$,*

$$\text{Ext}_{G(\mathbb{F}_q)}^i(M, N) \cong \text{Ext}_G^i(M, N \otimes \mathcal{G}_r(k)).$$

2.4 Good Filtrations. To make the desired computations of cohomology groups, we will make use of Proposition 2.3 (with $M = k = N$). In addition, we will use a special filtration on $\mathcal{G}_r(k)$. Recall that a G -module M has a *good filtration* if it admits a filtration with successive quotients of the form $H^0(\lambda)$, $\lambda \in X(T)_+$ [J, II 4.16].

One may consider $k[G]$ as a $G \times G$ -module via the left and right regular actions, respectively. A result due to Donkin and Koppinen [J, II 4.20] now says that $k[G]$ as a $G \times G$ -module admits a good filtration with factors of the form $H^0(\lambda) \otimes H^0(\lambda^*)$, each $\lambda \in X(T)_+$ appearing exactly once. Here $(g_1, g_2) \in G \times G$ acts via $g_1 \otimes g_2$ on each factor. If one takes the diagonal embedding of G into $G \times G$ one can use this fact to show that $k[G]$ admits a good filtration under the adjoint action of G .

For our purposes, we modify this slightly by using a partial Frobenius twist. Consider the composite

$$\phi : G \xrightarrow{\text{diag}} G \times G \xrightarrow{1 \times F^r} G \times G.$$

That is, when we take the diagonal embedding, we apply the Frobenius morphism r -times to the second factor giving $\phi(g) = (g, F^r(g))$. Let $G \times G$ act on $k[G]$ via the left and right regular representations as above, and then restrict this to a module over G via ϕ . Denote the resulting G -module by $k[G]^\vee$. The next proposition investigates filtrations on $k[G]^\vee$.

Proposition . *As G -modules $\mathcal{G}_r(k) \cong k[G]^\vee$. Moreover, $\mathcal{G}_r(k)$ has a filtration with factors of the form $H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}$ with multiplicity one for each $\lambda \in X(T)_+$.*

Proof. Let $f \in k[G]$ and $g, x \in G$. If we denote the action of G on $k[G]^\vee$ by \star then

$$(g \star f)(x) = f(g^{-1}x F^r(g)).$$

Next we define a Lang map $L : G \rightarrow G$ via $L(g) = g F^r(g^{-1})$. By setting $L^*(f) = f \circ L$ one obtains a bijection ([Hum2, 1.4])

$$L^* : k[G] \rightarrow \mathcal{G}_r(k) = \text{ind}_{G(\mathbb{F}_q)}^G(k) = \{f \in k[G] \mid f(gh) = f(g) \text{ for all } g \in G, h \in G(\mathbb{F}_q)\}.$$

Observe that

$$L^*(g \star f)(x) = (g \star f)(L(x)) = f(g^{-1}L(x)F^r(g))$$

while

$$\begin{aligned} (g(L^*(f)))(x) &= L^*(f)(g^{-1}x) = f(L(g^{-1}x)) = f(g^{-1}x F^r(x^{-1}g)) \\ &= f(g^{-1}x F^r(x^{-1})F^r(g)) = f(g^{-1}L(x)F^r(g)). \end{aligned}$$

Hence, L^* is a G -equivariant bijective map from $k[G]^\vee$ onto $\mathcal{G}_r(k)$.

Since $k[G]$ viewed as a $G \times G$ -module, has a good filtration with factors $H^0(\lambda) \otimes H^0(\lambda^*)$, and $\phi(g)$ acts on each factor via $g \otimes F^r(g)$, it follows that $k[G]^\vee$ has a filtration with factors of the form $H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}$ and multiplicity one for each $\lambda \in X(T)_+$, as claimed. \square

2.5 Given weights $\lambda, \mu \in X(T)$, recall that we say $\mu < \lambda$ if and only if $\lambda - \mu = \sum_{\alpha \in \Delta} c_\alpha \alpha$ for integers $c_\alpha \geq 0$. That is, $\lambda - \mu$ must lie in the positive root lattice. In addition we say two weights λ and μ are linked if there exists an element w of the affine Weyl group such that $\mu = w \cdot \lambda$. Note that two weights $(\lambda_1, \lambda_2), (\mu_1, \mu_2) \in X(T \times T)$ are linked for the algebraic group $G \times G$ if and only if the components are linked for G .

For each dominant weight λ we define two finite saturated sets, namely

$$\pi_{<\lambda} = \{\mu \in X(T)_+ \mid \mu < \lambda\} \text{ and } \pi_{\leq\lambda} = \{\mu \in X(T)_+ \mid \mu \leq \lambda\}.$$

According to [J, II A.15] the $G \times G$ -module $k[G]$ has two submodules $M_{<\lambda}$ and $M_{\leq\lambda}$, both admitting good filtrations with factors $H^0(\nu) \otimes H^0(\nu^*)$ where each $\nu \in \pi_{<\lambda}$ ($\pi_{\leq\lambda}$, respectively) appears exactly once. By $S_{<\lambda}$ and $S_{\leq\lambda}$ we denote the $G \times G$ -summands of $M_{<\lambda}$ and $M_{\leq\lambda}$, respectively, whose $G \times G$ -composition factors have highest weights contained in the same $G \times G$ -linkage class as (λ, λ^*) . Similarly we define the quotients $Q_{\not<\lambda} = k[G]/S_{<\lambda}$ and $Q_{\not\leq\lambda} = k[G]/S_{\leq\lambda}$.

The group G acts on these modules via the embedding ϕ . From Proposition 2.4 and [J, II 4.17] one obtains the following result.

Theorem . *For each $\lambda \in X(T)_+$, there exist short exact sequences of G -modules*

$$0 \rightarrow S_{<\lambda} \rightarrow \mathcal{G}_r(k) \rightarrow Q_{\not<\lambda} \rightarrow 0$$

and

$$0 \rightarrow S_{\leq\lambda} \rightarrow \mathcal{G}_r(k) \rightarrow Q_{\not\leq\lambda} \rightarrow 0$$

with the following properties:

- (a) $S_{<\lambda}$ ($S_{\leq\lambda}$) has a filtration with factors of the form $H^0(\nu) \otimes H^0(\nu^*)^{(r)}$ where $\nu < \lambda$ ($\nu \leq \lambda$) and ν is linked to λ .
- (b) $Q_{\not<\lambda}$ ($Q_{\not\leq\lambda}$) has a filtration with factors of the form $H^0(\nu) \otimes H^0(\nu^*)^{(r)}$ where $\nu \not< \lambda$ ($\nu \not\leq \lambda$) or ν is not linked to λ .
- (c) The multiplicity in all cases is one.

2.6 A Vanishing Criterion. The next proposition illustrates the usefulness of the existence of the filtrations in Proposition 2.4.

Proposition (A). *Let M, N be rational G -modules and $i \geq 0$. If $\text{Ext}_G^i(M, N \otimes H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) = 0$ for all $\lambda \in X(T)_+$, then $\text{Ext}_{G(\mathbb{F}_q)}^i(M, N) = 0$.*

Proof. By the assumption and the long exact sequence in cohomology associated to a short exact sequence, it follows from Proposition 2.4 that $\text{Ext}_G^i(M, N \otimes \mathcal{G}_r(k)) = 0$ for all i . The result now follows from Proposition 2.3. \square

In particular, in the special case of $M = k = N$, we get the following criterion for a vanishing range in cohomology. We will see later how to identify such an m for a given G .

Proposition (B). *Let m be the least positive integer such that there exists $\lambda \in X(T)_+$ with $H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \neq 0$. Then $H^i(G(\mathbb{F}_q), k) \cong H^i(G, \mathcal{G}_r(k)) = 0$ for $0 < i < m$.*

2.7 Non-vanishing. While the identification of an m satisfying Proposition 2.6(B) gives a vanishing range, it does not a priori follow that $H^m(G(\mathbb{F}_q), k) \neq 0$. In this section, we develop some conditions under which this conclusion could be made, as well as conditions under which one might be able to precisely identify the cohomology group. As in Proposition 2.6(A), we have the following.

Proposition (A). *Let $\lambda \in X(T)_+$ and $M \in \{S_{<\lambda}, S_{\leq\lambda}, Q_{\not\leq\lambda}, Q_{\not<\lambda}\}$. Suppose that*

$$H^i(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) = 0$$

for all $\nu \in X(T)_+$ which appear in the filtration for M . Then $H^i(G, M) = 0$.

The next proposition reduces the problem of showing the non-vanishing of $H^m(G, \mathcal{G}_r(k))$ (and hence of $H^m(G(\mathbb{F}_q), k)$) to showing non-vanishing for a submodule of $\mathcal{G}_r(k)$.

Proposition (B). *Let m be a positive integer. For any $\lambda \in X(T)_+$,*

- (i) *if $H^{m+1}(G, S_{<\lambda}) = 0$, then the map $H^m(G, \mathcal{G}_r(k)) \rightarrow H^m(G, Q_{\not\leq\lambda})$ is surjective;*
- (ii) *if $H^m(G, S_{<\lambda}) = 0$, then the map $H^m(G, \mathcal{G}_r(k)) \rightarrow H^m(G, Q_{\not\leq\lambda})$ is injective;*
- (iii) *if the conditions in (i) and (ii) hold, then $H^m(G, \mathcal{G}_r(k)) \cong H^m(G, Q_{\not\leq\lambda})$.*

Proof. Consider the short exact sequence

$$0 \rightarrow S_{<\lambda} \rightarrow \mathcal{G}_r(k) \rightarrow Q_{\not\leq\lambda} \rightarrow 0$$

and the associated long exact sequence in cohomology

$$\cdots \rightarrow H^m(G, S_{<\lambda}) \rightarrow H^m(G, \mathcal{G}_r(k)) \rightarrow H^m(G, Q_{\not\leq\lambda}) \rightarrow H^{m+1}(G, S_{<\lambda}) \rightarrow \cdots$$

The claims in (i) and (ii) follow immediately from this exact sequence. \square

Remark . In order to show the non-vanishing of $H^m(G(\mathbb{F}_q), k)$, condition (i) is the crucial condition. Whereas condition (ii) potentially allows us to identify $H^m(G(\mathbb{F}_q), k)$ precisely. Note that condition (ii) is immediately satisfied by any weight λ which is minimal with respect to the above standard ordering such that $H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \neq 0$.

Having reduced the problem to $H^i(G, Q_{\not\leq\lambda})$, we make some similar homological observations about this group.

Proposition (C). *Let m be the least positive integer such that there exists $\nu \in X(T)_+$ with $H^m(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) \neq 0$. For any $\lambda \in X(T)_+$,*

- (i) *the map $H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \rightarrow H^m(G, Q_{\not\leq\lambda})$ is injective;*
- (ii) *if $H^m(G, Q_{\not\leq\lambda}) = 0$, then $H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \cong H^m(G, Q_{\not\leq\lambda})$.*

Proof. Consider the short exact sequence

$$0 \rightarrow H^0(\lambda) \otimes H^0(\lambda^*)^{(r)} \rightarrow Q_{\not\leq\lambda} \rightarrow Q_{\not\leq\lambda} \rightarrow 0$$

and the associated LES

$$\cdots \rightarrow H^{m-1}(G, Q_{\not\leq\lambda}) \rightarrow H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \rightarrow H^m(G, Q_{\not\leq\lambda}) \rightarrow H^m(G, Q_{\not\leq\lambda}) \rightarrow \cdots$$

If $m > 1$, the claims follow immediately since the first term is zero by minimality of m .

Suppose that $m = 1$. Suppose first that λ is linked to the zero weight. Since we certainly cannot have $0 > \lambda$, by Theorem 2.5(b), $H^0(G, Q_{\not\prec\lambda}) = \text{Hom}_G(k, Q_{\not\prec\lambda}) = 0$, and the argument follows as above.

Suppose now that λ is not linked to the zero weight. The module $Q_{\not\prec\lambda}$ may be decomposed as a direct sum $M_1 \oplus M_2$ where M_1 has a filtration with factors of the form $H^0(\nu) \otimes H^0(\nu^*)^{(1)}$ with ν linked to λ and M_2 has such a filtration with ν not linked to λ . Then $H^1(G, Q_{\not\prec\lambda}) = H^1(G, M_1) \oplus H^1(G, M_2)$. Consider the short exact sequence

$$(2.7.1) \quad 0 \rightarrow H^0(\lambda) \otimes H^0(\lambda^*)^{(1)} \rightarrow M_1 \rightarrow M'_1 \rightarrow 0,$$

which defines a module M'_1 with a corresponding filtration. Observe that $Q_{\not\prec\lambda} \cong M'_1 \oplus M_2$, and hence $H^1(G, Q_{\not\prec\lambda}) \cong H^1(G, M'_1) \oplus H^1(G, M_2)$. Since λ is not linked to zero, $H^0(G, M'_1) = \text{Hom}_G(k, M'_1) = 0$. Arguing as above with the LES associated to (2.7.1) gives

$$H^1(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \hookrightarrow H^1(G, M_1) \hookrightarrow H^1(G, Q_{\not\prec\lambda}),$$

and so part (i) holds. For part (ii), it follows from the additional assumption that $H^1(G, M'_1) = 0$ and $H^1(G, M_2) = 0$. Again using the LES associated to (2.7.1), it follows that

$$H^1(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \cong H^1(G, M_1) = H^1(G, M_1) \oplus H^1(G, M_2) \cong H^1(G, Q_{\not\prec\lambda})$$

as claimed. □

2.8 Combining the propositions in the preceding section, we can obtain a condition under which we have sharp vanishing bounds, and an explicit identification of $H^m(G(\mathbb{F}_q), k)$ with a single G -cohomology group.

Theorem (A). *Let m be the least positive integer such that there exists $\nu \in X(T)_+$ with $H^m(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) \neq 0$. Let $\lambda \in X(T)_+$ be such that $H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \neq 0$. Suppose $H^{m+1}(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) = 0$ for all $\nu < \lambda$ that are linked to λ . Then*

- (i) $H^i(G(\mathbb{F}_q), k) = 0$ for $0 < i < m$;
- (ii) $H^m(G(\mathbb{F}_q), k) \neq 0$;
- (iii) *if, in addition, $H^m(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) = 0$ for all $\nu \in X(T)_+$ with $\nu \neq \lambda$, then $H^m(G(\mathbb{F}_q), k) \cong H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)})$.*

Proof. Part (i) follows from Proposition 2.6(B). For part (ii), note first that, by the hypothesis on λ and Proposition 2.7(C)(i), $H^m(G, Q_{\not\prec\lambda}) \neq 0$. On the other hand, by the hypothesis on weights less than λ and Proposition 2.7(A), $H^{m+1}(G, S_{<\lambda}) = 0$. Hence, by Proposition 2.7(B)(i), the map $H^m(G, \mathcal{G}_r(k)) \rightarrow H^m(G, Q_{\not\prec\lambda})$ is surjective. Therefore, by Proposition 2.3, $H^m(G(\mathbb{F}_q), k) \cong H^m(G, \mathcal{G}_r(k)) \neq 0$ as claimed.

For part (iii), by the added assumption and Proposition 2.7(A), we have $H^m(G, S_{<\lambda}) = 0$ and $H^m(G, Q_{\not\prec\lambda}) = 0$. By Proposition 2.7(B)(iii) and Proposition 2.7(C)(ii), we have $H^m(G, \mathcal{G}_r(k)) \cong H^m(G, Q_{\not\prec\lambda}) \cong H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)})$ as claimed. □

From the filtration on $\mathcal{G}_r(k)$ in Proposition 2.4, $H^i(G(\mathbb{F}_q), k) \cong H^i(G, \mathcal{G}_r(k))$ can be decomposed as a direct sum over linkage classes of dominant weights. As such, using an analogous argument, a slightly weaker condition for non-vanishing can be obtained.

Theorem (B). *For a fixed linkage class \mathcal{L} , let m be the least positive integer such that there exists $\nu \in \mathcal{L}$ with $H^m(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) \neq 0$. Let $\lambda \in \mathcal{L}$ be such that $H^m(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \neq 0$. Suppose $H^{m+1}(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)}) = 0$ for all $\nu < \lambda$ in \mathcal{L} . Then $H^m(G(\mathbb{F}_q), k) \neq 0$.*

3 Properties of the Cohomology Groups

In Section 2, it was shown that knowledge of cohomology groups of the form $H^i(G, H^0(\nu) \otimes H^0(\nu^*)^{(r)})$ for $\nu \in X(T)_+$ provides information on $H^i(G(\mathbb{F}_q), k)$. In this section, we study these G -cohomology groups and collect a number of useful properties that will be used throughout the remainder of the paper.

3.1 Reducing to G_r -cohomology. We will make frequent use of the following identification of G -extensions with G_r -cohomology.

Lemma . *Let $\nu_1, \nu_2 \in X(T)_+$. Assume that $H^j(G_r, H^0(\nu_1))^{(-r)}$ admits a good filtration for all $j > 0$. Then for all j*

$$H^j(G, H^0(\nu_1) \otimes H^0(\nu_2^*)^{(r)}) \cong \text{Ext}_G^j(V(\nu_2)^{(r)}, H^0(\nu_1)) \cong \text{Hom}_G(V(\nu_2), H^j(G_r, H^0(\nu_1))^{(-r)}).$$

Proof. The first isomorphism is immediate. For the crucial second isomorphism, consider the Lyndon-Hochschild-Serre spectral sequence

$$E_2^{i,j} = \text{Ext}_{G/G_r}^i(V(\nu_2)^{(r)}, H^j(G_r, H^0(\nu_1))) \Rightarrow \text{Ext}_G^{i+j}(V(\nu_2)^{(r)}, H^0(\nu_1)).$$

We have

$$\begin{aligned} E_2^{i,j} &= \text{Ext}_{G/G_r}^i(V(\nu_2)^{(r)}, H^j(G_r, H^0(\nu_1))) \\ &\cong \text{Ext}_G^i(V(\nu_2), H^j(G_r, H^0(\nu_1))^{(-r)}). \end{aligned}$$

For $\nu \in X(T)_+$, $i > 0$, and V a G -module which admits a good filtration, we have $\text{Ext}_G^i(V(\nu), V) = 0$ (cf. [J, Prop. II 4.16]). By the hypothesis, we conclude that $E_2^{i,j} = 0$ for all $i > 0$ and the spectral sequence collapses to a single vertical column. This implies that $E_2^{0,j} \cong \text{Ext}_G^j(V(\nu_2)^{(r)}, H^0(\nu_1))$ for all j . \square

The assumption that $H^j(G_r, H^0(\nu))^{(-r)}$ admits a good filtration is a long-standing conjecture of Donkin. For $p > h$ (the Coxeter number of the root system associated to G), this is known for $r = 1$ by results of Andersen-Jantzen [AJ] and Kumar-Lauritzen-Thomsen [KLT]. For arbitrary r , this is known only for all degrees in the case $G = SL_2$. When r is arbitrary and $i = 1$ (p arbitrary) or $i = 2$ ($p \geq 3$), Bendel-Nakano-Pillen verified the assumption by direct computation [BNP4, BNP7]. Wright [W] has recently verified the $p = 2$, $i = 2$ case.

We will apply Lemma 3.1 at several points in the $r = 1$ case for direct applications to $G(\mathbb{F}_p)$ as well as inductively for dealing with $G(\mathbb{F}_q)$. As such, we will generally assume for the remainder of the paper that $p > h$.

3.2 Dimensions for $r = 1$. From Lemma 3.1, to obtain information about $H^i(G, H^0(\nu) \otimes H^0(\nu^*)^{(1)})$ for $\nu \in X(T)_+$, it suffices to consider $\text{Hom}_G(V(\nu), H^i(G_1, H^0(\nu)^{(-1)}))$. It is well-known that, from block considerations, $H^i(G_1, H^0(\nu)) = 0$ unless $\nu = w \cdot 0 + p\mu$ for $w \in W$ and $\mu \in X(T)$. For $p > h$, from [AJ] and [KLT], we have

$$(3.2.1) \quad H^i(G_1, H^0(\nu))^{(-1)} = \begin{cases} \text{ind}_B^G(S^{\frac{i-\ell(w)}{2}}(\mathfrak{u}^*) \otimes \mu) & \text{if } \nu = w \cdot 0 + p\mu \\ 0 & \text{otherwise,} \end{cases}$$

where $\mathfrak{u} = \text{Lie}(U)$. Note also that, since $p > h$ and ν is dominant, μ must also be dominant.

For a weight ν and $n \geq 0$, let $P_n(\nu)$ denote the dimension of the ν -weight space of $S^n(\mathfrak{u}^*)$. Equivalently, for $n > 0$, $P_n(\nu)$ denotes the number of times that ν can be expressed as a sum of exactly n positive roots, while $P_0(0) = 1$. The function P_n is often referred to as *Kostant's Partition Function*. By using [AJ, 3.8], [KLT, Thm 2], Lemma 3.1, and (3.2.1), we can give an explicit formula for the dimension of $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)})$. Namely,

Proposition . *Assume $p > h$. Let $\lambda = p\mu + w \cdot 0 \in X(T)_+$. Then*

$$\dim H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = \sum_{u \in W} (-1)^{\ell(u)} P_{\frac{i-\ell(w)}{2}}(u \cdot \lambda - \mu).$$

3.3 From Proposition 2.3 and Proposition 2.4, we can now deduce the following upper bound on the dimensions of the cohomology groups $H^i(G(\mathbb{F}_p), k)$.

Theorem . *Assume $p > h$.*

$$\dim H^i(G(\mathbb{F}_p), k) \leq \sum_{\{w \in W \mid \ell(w) \equiv i \pmod{2}\}} \sum_{\mu \in X(T)_+} \sum_{u \in W} (-1)^{\ell(u)} P_{\frac{i-\ell(w)}{2}}(u \cdot (p\mu + w \cdot 0) - \mu).$$

3.4 Degree Bounds. From our discussion in Section 2, to find vanishing ranges for $H^\bullet(G(\mathbb{F}_p), k)$ (or $H^\bullet(G(\mathbb{F}_q), k)$ more generally), a first step is to try to identify the least positive i such that $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)})$ is non-zero.

Assume that $p > h$ and $\lambda \in X(T)_+$ with $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$ for some $i > 0$. From Lemma 3.1 and the discussion in Section 3.2, we know that $\lambda = p\mu + w \cdot 0$ for $w \in W$ and $\mu \in X(T)_+$. Observe that if $\mu = 0$, in order for λ to be dominant, we must have $\lambda = 0$. But, $H^i(G, k) = 0$ for $i > 0$. Since we are interested in cohomology in non-zero degrees, we may safely assume that $\lambda, \mu \neq 0$. Corollary 3.5 below gives a relationship between i and the weight λ . We first derive a more general relationship that will be useful in inductive arguments.

Proposition . *Assume that $p > h$. Let $\gamma_1, \gamma_2 \in X(T)_+$, both non-zero, such that $\gamma_j = p\delta_j + w_j \cdot 0$ with $\delta_j \in X(T)_+$ and $w_j \in W$ for $j = 1, 2$. Assume $\text{Ext}_G^i(V(\gamma_2)^{(1)}, H^0(\gamma_1)) \neq 0$.*

- (a) *For $\sigma \in \Phi^+$, $p\langle \delta_2, \sigma^\vee \rangle - \langle \delta_1, \sigma^\vee \rangle + \ell(w_1) + \langle w_2 \cdot 0, \sigma^\vee \rangle \leq i$.*
- (b) *Let $\tilde{\alpha}$ denote the longest root in Φ^+ . Then $p\langle \delta_2, \tilde{\alpha}^\vee \rangle - \langle \delta_1, \tilde{\alpha}^\vee \rangle + \ell(w_1) - \ell(w_2) - 1 \leq i$.*

Proof. By Lemma 3.1, (3.2.1), and Frobenius reciprocity, we have

$$\begin{aligned} \mathrm{Ext}_G^i(V(\gamma_2)^{(1)}, H^0(\gamma_1)) &\cong \mathrm{Hom}_G(V(\gamma_2), H^i(G_1, H^0(\gamma_1))^{(-1)}) \\ &\cong \mathrm{Hom}_G(V(\gamma_2), \mathrm{ind}_B^G(S^{\frac{i-\ell(w_1)}{2}}(\mathfrak{u}^*) \otimes \delta_1)) \\ &\cong \mathrm{Hom}_B(V(\gamma_2), S^{\frac{i-\ell(w_1)}{2}}(\mathfrak{u}^*) \otimes \delta_1). \end{aligned}$$

Since this is non-zero, $\gamma_2 = p\delta_2 + w_2 \cdot 0$ must be a weight of $S^{\frac{i-\ell(w_1)}{2}}(\mathfrak{u}^*) \otimes \delta_1$. In other words, $\gamma_2 - \delta_1 = p\delta_2 - \delta_1 + w_2 \cdot 0$ must be a weight of $S^{\frac{i-\ell(w_1)}{2}}(\mathfrak{u}^*)$.

The vector space \mathfrak{u}^* has a basis of root vectors corresponding to positive roots. So a homogeneous weight of $S^j(\mathfrak{u}^*)$ is a sum of j not necessarily distinct positive roots. Therefore, $\gamma_2 - \delta_1$ must be expressible as a sum of $\frac{i-\ell(w_1)}{2}$ positive roots. Of course, for any positive roots σ_1, σ_2 , we have $\langle \sigma_1, \sigma_2^\vee \rangle \leq 2$. Hence, for $\sigma \in \Phi^+$, we have

$$\langle \gamma_2 - \delta_1, \sigma^\vee \rangle \leq \frac{i - \ell(w_1)}{2} * 2 = i - \ell(w_1).$$

Substituting $\gamma_2 - \delta_1 = p\delta_2 - \delta_1 + w_2 \cdot 0$ gives

$$p\langle \delta_2, \sigma^\vee \rangle - \langle \delta_1, \sigma^\vee \rangle + \langle w_2 \cdot 0, \sigma^\vee \rangle \leq i - \ell(w_1).$$

Part (a) immediately follows.

For part (b), recall from Observation 2.2(B) that $-w_2 \cdot 0$ can be expressed uniquely as a sum of precisely $\ell(w_2)$ distinct positive roots. Since at most one of those roots can be $\tilde{\alpha}$, and $\langle \tilde{\alpha}, \tilde{\alpha}^\vee \rangle = 2$, by Observation 2.2(A), we have

$$\langle -w_2 \cdot 0, \tilde{\alpha}^\vee \rangle \leq (\ell(w_2) - 1) * 1 + 2 = \ell(w_2) + 1$$

which gives part (b). \square

3.5 As a special case of Proposition 3.4 we have the following result.

Corollary . *Assume that $p > h$. Let $\lambda = p\mu + w \cdot 0$ be a non-zero dominant weight with $0 \neq \mu \in X(T)_+$ and $w \in W$. Assume $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$.*

- (a) *For $\sigma \in \Phi^+$, $(p-1)\langle \mu, \sigma^\vee \rangle + \ell(w) + \langle w \cdot 0, \sigma^\vee \rangle \leq i$.*
- (b) *Let $\tilde{\alpha}$ denote the longest root in Φ^+ . Then $(p-1)\langle \mu, \tilde{\alpha}^\vee \rangle - 1 \leq i$.*

Proof. Parts (a) and (b) follow immediately from Proposition 3.4 by taking $\gamma_1 = \lambda = \gamma_2$. \square

In the corollary, since μ is a non-zero dominant weight, $\langle \mu, \tilde{\alpha}^\vee \rangle \geq 1$. Hence, we immediately have that $i \geq p - 2$. It follows from Proposition 2.6(B) that $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < p - 2$. This will follow as a special case of a more general result in the next section.

4 A Minimal Vanishing Range

4.1 In this section, we use the preceding techniques to determine a general vanishing range for $H^i(G(\mathbb{F}_q), k)$ for $p > h$. We begin with some further extension properties that will be used in the proof.

Lemma . *Assume that $p > h$ and $r > 1$. Let $\lambda, \mu \in X(T)_+$, both non-zero, and $i \geq 0$. If $\text{Ext}_G^i(V(\lambda)^{(r)}, H^0(\mu)) \neq 0$, then there exists a non-zero weight $\gamma \in X(T)_+$ and nonnegative integers k, l such that*

- (a) $i = k + l$,
- (b) $\text{Ext}_G^k(V(\lambda)^{(r-1)}, H^0(\gamma)) \neq 0$,
- (c) $\text{Ext}_G^l(V(\gamma)^{(1)}, H^0(\mu)) \neq 0$, and
- (d) $\gamma = p\delta + w \cdot 0$ for some $w \in W$ and non-zero $\delta \in X(T)_+$.

Proof. Consider the Lyndon-Hochschild-Serre spectral sequence

$$E_2^{k,l} = \text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^l(G_1, H^0(\mu))) \Rightarrow \text{Ext}_G^{k+l}(V(\lambda)^{(r)}, H^0(\mu)).$$

The assumptions imply that there exist nonnegative integers k, l with $k + l = i$ and $\text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^l(G_1, H^0(\mu))) \neq 0$. The G -module $H^l(G_1, H^0(\mu))^{(-1)}$ has a good filtration. Therefore, there exists a dominant weight γ with

$$(4.1.1) \quad \text{Hom}_{G/G_1}(V(\gamma)^{(1)}, H^l(G_1, H^0(\mu))) \cong \text{Hom}_G(V(\gamma), H^l(G_1, H^0(\mu))^{(-1)}) \neq 0$$

and

$$(4.1.2) \quad \text{Ext}_G^k(V(\lambda)^{(r-1)}, H^0(\gamma)) \cong \text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^0(\gamma)^{(1)}) \neq 0.$$

Now (4.1.2) implies that γ is of the form $\gamma = p\delta + w \cdot 0$ with $w \in W$ and $\delta \in X(T)_+$. By Lemma 3.1 and (4.1.1), $\text{Ext}_G^l(V(\gamma)^{(1)}, H^0(\mu)) \cong \text{Hom}_G(V(\gamma), H^l(G_1, H^0(\mu))^{(-1)}) \neq 0$. This forces $\delta \neq 0$. Note that the assumptions also force $\mu = p\delta' + w' \cdot 0$ for some $w' \in W$ and nonzero $\delta' \in X(T)_+$. \square

4.2 Applying the lemma repeatedly immediately implies the following proposition.

Proposition . *Assume that $p > h$. Let $\lambda, \mu \in X(T)_+$, both non-zero, and $i \geq 0$. If $\text{Ext}_G^i(V(\lambda)^{(r)}, H^0(\mu)) \neq 0$, then there exists a sequence of non-zero weights $\mu = \gamma_0, \gamma_1, \dots, \gamma_{r-1}, \gamma_r = \lambda \in X(T)_+$ and nonnegative integers l_1, l_2, \dots, l_r such that*

- (a) $i = \sum_{j=1}^r l_j$,
- (b) $\text{Ext}_G^{l_j}(V(\gamma_j)^{(1)}, H^0(\gamma_{j-1})) \neq 0$, for $1 \leq j \leq r$ and
- (c) $\gamma_j = p\delta_j + u_j \cdot 0$ for some $u_j \in W$ and nonzero $\delta_j \in X(T)_+$.

4.3 The next step in our analysis is to obtain tighter control over a lower bound on i as in Proposition 4.2.

Proposition . *Assume that $p > h$. Let $0 \neq \lambda \in X(T)_+$ and $i \geq 0$. If $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \neq 0$, then there exists a sequence of non-zero weights $\lambda = \gamma_0, \gamma_1, \dots, \gamma_{r-1}, \gamma_r = \lambda \in X(T)_+$ such that $\gamma_j = p\delta_j + u_j \cdot 0$ for some $u_j \in W$ and nonzero $\delta_j \in X(T)_+$. Furthermore,*

$$(4.3.1) \quad i \geq \left(\sum_{j=1}^r (p-1) \langle \delta_j, \tilde{\alpha}^\vee \rangle \right) - r.$$

Proof. By Proposition 4.2 with $\lambda = \mu$, there exists a sequence of non-zero dominant weights $\lambda = \gamma_0, \gamma_1, \dots, \gamma_{r-1}, \gamma_r = \lambda$ with $\gamma_j = p\delta_j + u_j \cdot 0$ and corresponding nonnegative integers l_j with $i = \sum_{j=1}^r l_j$ and $\text{Ext}_G^{l_j}(V(\gamma_j)^{(1)}, H^0(\gamma_{j-1})) \neq 0$. The first part of the statement now follows.

For second part, we use Proposition 3.4(b) to obtain the inequalities

$$p\langle \delta_j, \tilde{\alpha}^\vee \rangle - \langle \delta_{j-1}, \tilde{\alpha}^\vee \rangle + \ell(u_{j-1}) - \ell(u_j) - 1 \leq l_j \text{ for } 1 \leq j \leq r.$$

Note that $\delta_0 = \delta_r$ and $u_0 = u_r$. Summing over j yields

$$\left(\sum_{j=1}^r (p-1)\langle \delta_j, \tilde{\alpha}^\vee \rangle \right) - r \leq \sum_{j=1}^r l_j = i.$$

□

4.4 For $p > h$ we can now present general vanishing ranges which address (1.1.1).

Theorem . *Assume that $p > h$. Then*

- (a) $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) = 0$ for $0 < i < r(p-2)$ and $\lambda \in X(T)_+$;
- (b) $H^i(G(\mathbb{F}_q), k) = 0$ for $0 < i < r(p-2)$.

Proof. Part (a) implies part (b) via Proposition 2.6(B). Suppose that

$$H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(r)}) \neq 0$$

for some $0 < i$ and $\lambda \in X(T)_+$. Clearly $\lambda \neq 0$, so we may apply Proposition 4.3. Since $\gamma_j \neq 0$, $1 \leq \langle \delta_j, \tilde{\alpha}^\vee \rangle$. Proposition 4.3 then gives $i \geq r(p-1) - r = r(p-2)$ as claimed. □

Observe that this vanishing range is generally larger than the one obtained in [H]. Precisely, the ranges obtained in [H] are of the form $0 < i < m$ where m depends on the root system. Except in certain type A_n cases, $m \leq r(p-1)/2$.

In the remainder of the paper, we further investigate this question to determine sharp bounds for root systems of type C_n (for all r ; see Theorem 5.4) and A_n (for $r = 1$ and generically for all r ; see Theorems 6.13, 6.14). In type C_n , the above bounds are in fact sharp.

Remark . Note that the assumption $\text{Ext}_G^i(V(\lambda)^{(r)}, H^0(\mu)) \neq 0$ in Proposition 4.2 can be replaced by $\text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^l(G_1, H^0(\mu))) \neq 0$, where $k+l = i$. In that case one arrives at the same conclusions with $l_1 = l$. Now the arguments used to prove Proposition 4.3 and Theorem 4.4 can be used to show that $\text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^l(G_1, H^0(\lambda))) = 0$ for all $k+l < r(p-2)$.

5 Type C_n , $n \geq 1$

Assume throughout this section that Φ is of type C_n , $n \geq 1$, and $p > h = 2n$.

5.1 Realization for $r = 1$. We determine the least $i > 0$ such that $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$. From Theorem 4.4, we know that $i \geq p-2$. Let $\tilde{\alpha} = 2\omega_1$ denote the longest positive root. We next construct a weight λ with $H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$.

Let $w = s_1 s_2 \dots s_{n-1} s_n s_{n-1} \dots s_2 s_1 \in W$. Then $-w \cdot 0 = n\tilde{\alpha} = 2n\omega_1$. Furthermore, when expressed as a sum of distinct positive roots, $-w \cdot 0$ consists of precisely all positive roots which contain an α_1 . Set $\lambda = p\omega_1 + w \cdot 0 = p\omega_1 - 2n\omega_1 = (p - 2n)\omega_1$. Then

$$\lambda - \omega_1 = (p - 1 - 2n)\omega_1 = \left(\frac{p-1}{2} - n \right) \tilde{\alpha}$$

is a highest weight of $S^j(\mathfrak{u}^*)$ where $j = \frac{p-1}{2} - n$.

We will apply Proposition 3.2 to compute $\dim \mathbb{H}^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)})$. Specifically, we will show that for $u \in W$

$$P_{\frac{p-1}{2}-n}(u \cdot \lambda - \omega_1) = \begin{cases} 1 & \text{if } u = 1 \\ 0 & \text{else.} \end{cases}$$

We will work with the ϵ -basis of $X(T)$. Rewrite $u \cdot \lambda - \omega_1 = ((p - 2n)u - 1)\epsilon_1 + u \cdot 0$ as $\sum_i c_i \epsilon_i$. In order for this expression to be a sum of positive roots, the coefficient c_1 has to be nonnegative. This forces $u(\epsilon_1) = \epsilon_1$. Then $u \cdot 0$ is of the form $-\sum_i d_i \alpha_i$ with $d_1 = 0$. This implies that $u \cdot \lambda - \omega_1 = (\frac{p-1}{2} - n)\tilde{\alpha} - \sum_i d_i \alpha_i$. Such an expression contains $p - 1 - 2n$ copies of α_1 . Since $\tilde{\alpha}$ is the only positive root containing $2\alpha_1$, the above expression can be written as a sum of $\frac{p-1}{2} - n$ positive roots if and only if $u = 1$. From Proposition 3.2, one concludes that $\mathbb{H}^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \cong k$.

5.2 More vanishing in degree $p - 2$. To get a precise vanishing range, we need to consider whether there are any other weights with non-zero cohomology in degree $p - 2$. Let $\lambda = p\mu + w \cdot 0 \in X(T)_+$ with $\mathbb{H}^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(-1)}) \neq 0$ for some $i > 0$. Consider the maximal short root $\alpha_0 = \omega_2$. By Corollary 3.5(a) with $\sigma = \alpha_0$,

$$(5.2.1) \quad i \geq (p - 1)\langle \mu, \alpha_0^\vee \rangle + \ell(w) + \langle w \cdot 0, \alpha_0^\vee \rangle.$$

There are three positive roots β with $\langle \beta, \alpha_0^\vee \rangle = 2$ (unless $n = 1$, in which case there is only one and $\alpha_0 = 2\omega_1$). Since $-w \cdot 0$ can be expressed uniquely as a sum of $\ell(w)$ distinct positive roots, we can say $\langle -w \cdot 0, \alpha_0^\vee \rangle \leq \ell(w) + 3$. Hence, (5.2.1) can be rewritten as

$$(5.2.2) \quad i \geq (p - 1)\langle \mu, \alpha_0^\vee \rangle - 3.$$

In Type C_n , $\langle \omega_1, \alpha_0^\vee \rangle = 1$. But, for $2 \leq j \leq n$, $\langle \omega_j, \alpha_0^\vee \rangle = 2$. Hence, for $\mu \in X(T)_+$, if $\mu \neq 0, \omega_1$, then $\langle \mu, \alpha_0^\vee \rangle \geq 2$. If $\langle \mu, \alpha_0^\vee \rangle \geq 2$, then (5.2.2) becomes

$$i \geq 2(p - 1) - 3 = 2p - 5 > p - 2$$

since $p \geq 5$ ($n \geq 2$). Therefore, the only candidates for a non-zero cohomology group in degree $p - 2$ are with $\lambda = p\omega_1 + w \cdot 0$ for some $w \in W$. This makes sense because the weight constructed in Section 5.1 is of this form.

5.3 A sharp bound for $r = 1$. Suppose $\lambda = p\omega_1 + w \cdot 0 \in X(T)_+$ and

$$\mathbb{H}^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0.$$

As in the proof of Proposition 3.4, $\lambda - \omega_1 = (p - 1)\omega_1 + w \cdot 0$ must be a weight of $S^{\frac{p-2-\ell(w)}{2}}(\mathfrak{u}^*)$. Also, by Corollary 3.5, we must have $\langle w \cdot 0, \tilde{\alpha}^\vee \rangle = -(\ell(w) + 1)$. Recall that $\tilde{\alpha} = 2\omega_1$ and $\langle \lambda - \omega_1, \tilde{\alpha}^\vee \rangle$ counts the number of times that α_1 appears in $\lambda - \omega_1$. Therefore, $(p - 1)\omega_1 + w \cdot 0$

contains exactly $(p-1) - (\ell(w) + 1) = p - \ell(w) - 2$ copies of α_1 . To obtain that many α_1 s as sum of $\frac{p-\ell(w)-2}{2}$ positive roots requires each root to contain two α_1 s, i.e., all roots are equal to $\tilde{\alpha}$ and $-w \cdot 0 = (\ell(w) + 1)\tilde{\alpha}$. Since $-w \cdot 0$ can be expressed in a unique way as a sum of distinct roots, one concludes that $-w \cdot 0$ is actually the sum of all roots that contain α_1 . Hence, $\lambda = (p-2n)\omega_1$, the weight given in Section 5.1. So the λ exhibited there is the only dominant weight with $H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$.

Note that $\lambda = (p-2n)\omega_1$ is contained in the lowest alcove. There are no smaller dominant weights that are linked to λ (so the condition in Theorem 2.8(A) involving H^{m+1} is vacuous). Consequently, Theorem 2.8(A) and the above discussion now yields:

Theorem . *Suppose Φ is of type C_n with $p > 2n$. Then*

- (a) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < p-2$;
- (b) $H^{p-2}(G(\mathbb{F}_p), k) \cong k$.

5.4 A sharp bound for all r . In this section we will address (1.1.1) and (1.1.2) in general for $H^\bullet(G(\mathbb{F}_q), k)$ when Φ is of type C_n .

Lemma . *Assume $p > 2n$. Let $\lambda = (p-2n)\omega_1$. Then $\text{Ext}_G^{r(p-2)}(V(\lambda)^{(r)}, H^0(\lambda)) \neq 0$.*

Proof. We use induction on r . If $r = 1$ the assertion follows from Section 5.1. Next we make use of the LHS spectral sequence

$$E_2^{k,l} = \text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^l(G_1, H^0(\lambda))) \Rightarrow \text{Ext}_G^{k+l}(V(\lambda)^{(r)}, H^0(\lambda)).$$

Our discussion in Section 5.1 implies that

$$\text{Hom}_G(V(\lambda), H^l(G_1, H^0(\lambda))^{(-1)}) \cong \text{Hom}_B(V(\lambda), S^{\frac{l-2n+1}{2}}(\mathfrak{u}^*) \otimes \omega_1) \cong \begin{cases} k & \text{if } l = p-2, \\ 0 & \text{if } l < p-2. \end{cases}$$

Moreover, for $j \leq \frac{p-1}{2} - n$, all weights γ appearing in $S^j(\mathfrak{u}^*) \otimes \omega_1$ satisfy $\langle \gamma + \rho, \alpha_0^\vee \rangle < p$. Hence, $H^l(G_1, H^0(\lambda))^{(-1)}$ is completely reducible for $l \leq p-2$. In addition, $H^0(\lambda) = L(\lambda)$ appears as a summand if and only if $l = p-2$. Clearly the trivial module does not appear as a summand of $H^l(G_1, H^0(\lambda))^{(-1)}$. The linkage principle now forces $E_2^{k,l} = 0$ whenever $l < p-2$. Further, writing $H^l(G_1, H^0(\lambda))^{(-1)} \cong H^0(\lambda) \oplus N$ for some module N ,

$$\begin{aligned} E_2^{(r-1)(p-2), p-2} &= \text{Ext}_{G/G_1}^{(r-1)(p-2), p-2}(V(\lambda)^{(r)}, H^{p-2}(G_1, H^0(\lambda))) \\ &\cong \text{Ext}_G^{(r-1)(p-2)}(V(\lambda)^{(r-1)}, H^{p-2}(G_1, H^0(\lambda))^{-1}) \\ &= \text{Ext}_G^{(r-1)(p-2)}(V(\lambda)^{(r-1)}, H^0(\lambda) \oplus N) \\ &\cong \text{Ext}_G^{(r-1)(p-2)}(V(\lambda)^{(r-1)}, H^0(\lambda)) \oplus \text{Ext}_G^{(r-1)(p-2)}(V(\lambda)^{(r-1)}, N). \end{aligned}$$

By induction, the first summand in this last line is non-zero, and hence $E_2^{(r-1)(p-2), p-2} \neq 0$. Finally, the remark in Section 4.4 says that $E_2^{k,l} = 0$ whenever $k+l < r(p-2)$. Therefore, the $((r-1)(p-2), p-2)$ -term of the E_2 -page transgresses to the E_∞ -page which produces a non-trivial cohomology class in degree $r(p-2)$. \square

By applying Theorem 4.4, the fact that λ is the smallest weight in its linkage class, and Theorem 2.8(B) one obtains the following theorem.

Theorem . *Suppose Φ is of type C_n with $p > 2n$. Then*

- (a) $H^i(G(\mathbb{F}_q), k) = 0$ for $0 < i < r(p-2)$;
- (b) $H^{r(p-2)}(G(\mathbb{F}_q), k) \neq 0$.

6 Type A_n , $n \geq 2$

Assume throughout this section that Φ is of type A_n , $n \geq 2$, and that $p > h = n + 1$. Note that type A_1 is equivalent to type C_1 which was covered in Section 5.

6.1 An upper bound for $r = 1$. We first construct a weight λ with

$$H^{2p-3}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0.$$

Set $w := s_1 s_2 \dots s_{n-1} s_n s_{n-1} \dots s_2 s_1 \in W$, where s_i is again the reflection corresponding to the i th simple root α_i . Then $\ell(w) = 2n - 1$ and $-w \cdot 0 = n\tilde{\alpha}$. When decomposed uniquely into a sum of distinct positive roots, $-w \cdot 0$ consists of precisely all positive roots which contain either an α_1 or an α_n (or both). Set $\lambda := p\tilde{\alpha} + w \cdot 0 = (p-n)\tilde{\alpha}$ and $\mu := \tilde{\alpha}$. Then $\lambda - \mu = (p-n-1)\tilde{\alpha}$ is a weight of $S^{p-n-1}(\mathfrak{u}^*)$. Indeed, it is the highest weight corresponding to taking $(p-n-1)$ -copies of $\phi_{\tilde{\alpha}} \in \mathfrak{u}^*$ (the root vector corresponding to $\tilde{\alpha}$). Similar to the argument in Section 5.1, we will show that

$$P_{p-n-1}(u \cdot \lambda - \mu) = \begin{cases} 1 & \text{if } u = 1 \\ 0 & \text{else.} \end{cases}$$

We will work with the ϵ -basis of $X(T)$. Rewrite $u \cdot \lambda - \mu = ((p-n)u - 1)(\epsilon_1 - \epsilon_{n+1}) + u \cdot 0$ as $\sum_i c_i \epsilon_i$. In order for this expression to be a sum of positive roots, the coefficient c_1 has to be nonnegative and c_{n+1} has to be less than or equal to zero. This forces $u(\epsilon_1) = \epsilon_1$ and $u(\epsilon_{n+1}) = \epsilon_{n+1}$. This forces now $u \cdot 0$ to be of the form $-\sum_i d_i \alpha_i$ with $d_1 = d_{n+1} = 0$. This implies that $u \cdot \lambda - \mu = (p-n-1)\tilde{\alpha} - \sum_i d_i \alpha_i$ can be written as a sum of $p-n-1$ positive roots if and only if $u = 1$.

Proposition . *Suppose Φ is of type A_n with $n \geq 2$ and $p \geq n + 2$. Let $\lambda = (p-n)\tilde{\alpha} = (p-n)\omega_1 + (p-n)\omega_n$. Then*

- (a) $H^{2p-3}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = k$;
- (b) $H^{2p-3}(G(\mathbb{F}_p), k) \neq 0$.

Proof. Part (a) follows from our analysis above and Proposition 3.2.

For part (b) suppose that $0 \neq \mu \in X(T)_+$ is linked to $(p-n)\tilde{\alpha}$ and $H^i(G, H^0(\mu) \otimes H^0(\mu^*)^{(1)}) \neq 0$ for some $i > 0$. As noted in Section 3.2, we necessarily have $\mu = p\delta + w \cdot 0$ for some $0 \neq \delta \in X(T)_+$ and $w \in W$. Observe that, since μ lies in the root lattice, δ also lies in the root lattice. Therefore, $\langle \delta, \tilde{\alpha}^\vee \rangle \geq 2$. From Corollary 3.5(b) we get $i \geq 2(p-1) - 1 = 2p - 3$.

Note also that $(p-n)\tilde{\alpha}$ lies in the second fundamental p -alcove. Indeed, it is the reflection of the zero weight across the upper wall. So the only weight μ of the form $p\tilde{\alpha} + w \cdot 0$ with

$\mu < \lambda$ would be the zero weight. And we know that $H^i(G, k) = 0$ for all $i > 0$. Therefore, we can apply Theorem 2.8(B) to deduce the result. \square

6.2 An upper bound for $r > 1$. The following result identifies a non-zero cohomology class in degree $r(2p - 3)$.

Proposition . *Suppose Φ is of type A_n with $n \geq 2$ and $p \geq n + 2$. Let $\lambda = (p - n)\tilde{\alpha} = (p - n)\omega_1 + (p - n)\omega_n$ and γ be a weight that is linked to the zero weight. Then*

$$(a) \text{Ext}_G^i(V(\gamma)^{(1)}, H^0(\lambda)) \cong \text{Hom}_G(V(\gamma), H^i(G_1, H^0(\lambda))^{(-1)}) \cong \begin{cases} k & \text{if } i = 2p - 3 \\ & \text{and } \gamma = \lambda, \\ 0 & \text{if } i < 2p - 3; \end{cases}$$

$$(b) H^{r(2p-3)}(G(\mathbb{F}_q), k) \neq 0.$$

Proof. (a) Note that $\lambda = (p - n)\tilde{\alpha}$ does not appear as a weight in $S^i(\mathfrak{u}^*) \otimes \tilde{\alpha}$ if $i < p - n - 1$. The discussion in the previous section and Lemma 3.1 yields part (a).

(b) We will now follow the argument of Section 5.4 for type A and the weight $\lambda = (p - n)\tilde{\alpha} = (p - n)\omega_1 + (p - n)\omega_n$ to show that $\text{Ext}_G^{r(2p-3)}(V(\lambda)^{(r)}, H^0(\lambda)) \neq 0$.

Part (a) implies that the $E_2^{k,l}$ -terms of the LHS spectral sequence

$$E_2^{k,l} = \text{Ext}_{G/G_1}^k(V(\lambda)^{(r)}, H^l(G_1, H^0(\lambda))) \Rightarrow \text{Ext}_G^{k+l}(V(\lambda)^{(r)}, H^0(\lambda))$$

vanish for $l < 2p - 3$. Moreover, $H^0(\lambda)$ is a summand of $H^{2p-3}(G_1, H^0(\lambda))^{(-1)}$ and by using induction on r one obtains that $E_2^{(r-1)(2p-3), 2p-3} \neq 0$.

Note that if we apply Proposition 4.3 for our specific weight λ all elements in the sequence $\gamma_0, \dots, \gamma_r$ have to be non-zero, dominant, and linked to zero. This implies that $\gamma_j = p\delta_j + u_j \cdot 0$ with $2 \leq \langle \delta_j, \tilde{\alpha}^\vee \rangle$. Therefore, equation (4.3.1) now reads

$$r(2p - 3) \leq \sum_{j=1}^r (p - 1) \langle \delta_j, \tilde{\alpha}^\vee \rangle - r \leq \sum_{j=1}^r l_j = i.$$

As in Section 5.4 we use the remark of Section 4.4 and conclude that $E_2^{k,l} = 0$ whenever $k + l < r(2p - 3)$. Hence,

$$k \cong \text{Ext}_{G/G_1}^{(r-1)(2p-3)}(V(\lambda)^{(r)}, H^{2p-3}(G_1, H^0(\lambda))) \hookrightarrow \text{Ext}_G^{r(2p-3)}(V(\lambda)^{(r)}, H^0(\lambda)).$$

Part (b) now follows because $\lambda = (p - n)\tilde{\alpha}$ is the lowest non-zero weight in its linkage class. \square

Proposition 6.2 and Theorem 4.4 imply that the least positive i with $H^i(G(\mathbb{F}_q), k) \neq 0$ satisfies $r(p - 2) \leq i \leq r(2p - 3)$. For $r = 1$, in the following sections, we identify precisely the value of i . The answer will depend on the relationship between p and n .

6.3 Counting simple roots. Let $\alpha_1, \alpha_2, \dots, \alpha_n$ denote the simple roots and $\omega_1, \omega_2, \dots, \omega_n$ the corresponding fundamental weights. Any weight γ can be written in the form $\gamma = \sum_{j=1}^n c_j \alpha_j$ with $c_j \in \mathbb{Q}$. We define $M_j(\gamma) := c_j$, $M(\gamma) := \max\{c_j\}$ and $m(\gamma) := \max\{j \mid c_j = M(\gamma)\}$. In addition we set $N_j := j(n + 1 - j)$. We make the following

Observation . For $1 \leq j \leq n$ and $w \in W$,

- (a) $\omega_j = \frac{1}{n+1}$ (sum of all positive roots which contain α_j),
- (b) N_j is the number of positive roots in Φ which contain α_j ,
- (c) $m(\omega_j) = j$,
- (d) $M_{m(\omega_j)}(\omega_j) = M_j(\omega_j) = M(\omega_j) = \frac{N_j}{n+1}$,
- (e) $M_j(2\rho) = N_j$,
- (f) $M_j(2\rho) = M_j(-w_0 \cdot 0) \geq M_j(-w \cdot 0)$.

Suppose we have dominant weights λ, μ with $\lambda = p\delta_2 + w_2 \cdot 0$, $\mu = p\delta_1 + w_1 \cdot 0$ and $H^i(G, H^0(\mu) \otimes H^0(\lambda^*)^{(1)}) \neq 0$ for some $i > 0$. As in the proof of Proposition 3.4, $\lambda - \delta_1$ must be a weight of $S^{\frac{i-\ell(w_1)}{2}}(\mathfrak{u}^*)$. Hence $M_{m(\delta_2)}(\lambda - \delta_1) \leq \frac{i-\ell(w_1)}{2}$. Using $\lambda = p\delta_2 + w_2 \cdot 0$, it follows that

$$(6.3.1) \quad 2pM_{m(\delta_2)}(\delta_2) + 2M_{m(\delta_2)}(w_2 \cdot 0) - 2M_{m(\delta_2)}(\delta_1) + l(w_1) \leq i.$$

Note that $M_{m(\delta_2)}(\delta_1) \leq M(\delta_1) = M_{m(\delta_1)}(\delta_1)$ and that $-M_{m(\delta_1)}(w_1 \cdot 0) \leq l(w_1)$ (from Observation 2.2(B)). One obtains

$$(6.3.2) \quad 2pM_{m(\delta_2)}(\delta_2) + 2M_{m(\delta_2)}(w_2 \cdot 0) - 2M_{m(\delta_1)}(\delta_1) - M_{m(\delta_1)}(w_1 \cdot 0) \leq i.$$

Next assume that $\mu = \lambda$. Then (6.3.2) results in $2(p-1)M_{m(\delta_1)}(\delta_1) + M_{m(\delta_1)}(w_1 \cdot 0) \leq i$. Using Observation (f) gives

$$(6.3.3) \quad 2(p-1)M_{m(\delta_1)}(\delta_1) + M_{m(\delta_1)}(-2\rho) \leq i.$$

Suppose now that $\delta_1 = \omega_j$ for $1 \leq j \leq n$. From Observations (d) and (e), (6.3.3) becomes

$$(6.3.4) \quad 2 \left(\frac{p-1}{n+1} \right) N_j - N_j \leq i.$$

However, we can say more than this. Suppose that in fact $M_j(-w_1 \cdot 0) = N_j$. Then, when expressed as a sum of distinct positive roots, $-w_1 \cdot 0$ contains all N_j roots containing α_j and possibly some other positive roots. In other words, $-w_1 \cdot 0 = (n+1)\omega_j + \sigma$ where σ is a sum of distinct positive roots not containing α_j . Then

$$\lambda = p\omega_j + w_1 \cdot 0 = p\omega_j - (n+1)\omega_j - \sigma = (p-n-1)\omega_j - \sigma \leq (p-n-1)\omega_j.$$

Hence, the only way λ can be dominant is if $\sigma = 0$. In other words, $\lambda = (p-n-1)\omega_j$ and we have shown the following.

Proposition . Suppose that Φ is of type A_n with $n \geq 2$ and $\lambda = p\omega_j + w \cdot 0 \in X(T)_+$ for $w \in W$ is a weight of $S^{\frac{i-\ell(w)}{2}}(\mathfrak{u}^*) \otimes \omega_j$. Then

$$i \geq \left[2 \left(\frac{p-1}{n+1} \right) - 1 \right] j(n+1-j)$$

with equality possible if and only if $\lambda = (p-n-1)\omega_j$.

Remark . The assumption that $\lambda = p\omega_j + w \cdot 0$ is a weight of $S^{\frac{i-\ell(w)}{2}}(\mathfrak{u}^*) \otimes \omega_j$ places restrictions on p (and n). Indeed, $\lambda - \omega_j = (p-1)\omega_j + w \cdot 0$ must lie in the root lattice. But $w \cdot 0$ lies in the root lattice. Therefore, $(p-1)\omega_j$ must also lie in the root lattice. However,

for $1 \leq j \leq (n+1)/2$, to have $(p-1)\omega_j$ or symmetrically $(p-1)\omega_{n+1-j}$ in the root lattice, we must have that $(n+1)$ divides $(p-1)j$.

Given the constraints noted in the remark, it is useful to rewrite the right hand side of the proposition as

$$(6.3.5) \quad \left[2 \left(\frac{p-1}{n+1}\right) - 1\right] j(n+1-j) = [2(p-1) - (n+1)]j \left(1 - \frac{j}{n+1}\right).$$

6.4 Larger Weights. In this section, we will see that the only non-zero non-fundamental dominant weight λ which can have $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$ for $0 \leq i \leq 2p-3$ is the weight $\lambda = (p-n)\tilde{\alpha}$ considered in Section 6.1. Indeed, observe that when $p \geq n+2$, $2(p-1) + 2(n-1) \left[\frac{p-1}{n+1} - 1\right] > 2p-3$.

Proposition . *Suppose that Φ is of type A_n with $n \geq 2$ and $p \geq n+2$. Let $\lambda = p\mu + w \cdot 0 \in X(T)_+$ for $w \in W$ with $\langle \mu, \tilde{\alpha} \rangle \geq 2$. If $\lambda \neq (p-n)\tilde{\alpha}$, then $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for*

$$0 < i \leq 2(p-1) + 2(n-1) \left[\frac{p-1}{n+1} - 1\right].$$

Proof. Let λ and μ be as given. Assume that $i \neq 0$ and $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$. Using our definition of $m(\mu)$ from Section 6.3 we write $\mu = a\omega_{m(\mu)} + \sigma$, where σ is a sum of fundamental weights other than $\omega_{m(\mu)}$ and a is a positive integer. Note that σ is nonzero if $a = 1$. Set $j = \min\{m(\mu), n+1 - m(\mu)\}$. Note that $M_{m(\mu)}(\omega_l) \geq \frac{j}{n+1}$ for any $l \neq j$. We obtain the following inequality

$$M_{m(\mu)}(\mu) \geq \begin{cases} \frac{2N_j}{n+1} & \text{if } a \geq 2, \\ \frac{N_j+j}{n+1} & \text{else.} \end{cases}$$

Substituting the above into equation (6.3.3) with $\delta_1 = \mu$ yields

$$i \geq \begin{cases} 2 \left(\frac{p-1}{n+1}\right) (2N_j) - N_j & \text{if } a \geq 2, \\ 2 \left(\frac{p-1}{n+1}\right) (N_j + j) - N_j & \text{else.} \end{cases}$$

As a function of j , N_j is increasing on the interval $(0, (n+1)/2)$. Therefore both of the above expressions are minimal when j is as small as possible. In the first case $j = 1$ is possible. However, since we are assuming that $\mu \neq \omega_1 + \omega_n$, we may assume that $j \geq 2$ for the second case. One obtains

$$(6.4.1) \quad i \geq \begin{cases} 2 \left(\frac{p-1}{n+1}\right) 2n - n & \text{if } a \geq 2, \\ 2 \left(\frac{p-1}{n+1}\right) (2(n-1) + 2) - 2(n-1) = 2 \left(\frac{p-1}{n+1}\right) 2n - 2(n-1) & \text{else.} \end{cases}$$

Hence,

$$\begin{aligned}
 i &\geq 4n \binom{p-1}{n+1} - 2(n-1) \\
 &= 4(n+1) \binom{p-1}{n+1} - 4 \binom{p-1}{n+1} - 2(n-1) \\
 &= 2(p-1) + [2(n+1) - 4] \binom{p-1}{n+1} - 2(n-1) \\
 &= 2(p-1) + 2(n-1) \binom{p-1}{n+1} - 2(n-1) \\
 &= 2(p-1) + 2(n-1) \left[\frac{p-1}{n+1} - 1 \right].
 \end{aligned}$$

□

To determine sharp vanishing bounds, we need to consider the relationship between p and n . This will be done in the succeeding sections.

6.5 The Case: $\gcd(p-1, n+1) = 1$. It follows from Remark 6.3 that under this assumption, the weight $(p-1)\omega_j$ does not lie in the root lattice for any j . Therefore, μ must be the sum of at least two (not necessarily distinct) fundamental dominant weights, and $\langle \mu, \tilde{\alpha}^\vee \rangle \geq 2$. From Corollary 3.5(b), we conclude that $i \geq 2p-3$. Combining this with Proposition 6.1, Proposition 6.4, and Theorem 2.8(A), we obtain these sharp bounds.

Theorem . *Suppose Φ is of type A_n with $n \geq 2$, $p \geq n+2$ and $\gcd(p-1, n+1) = 1$. Then*

- (a) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p-3$;
- (b) $H^{2p-3}(G(\mathbb{F}_p), k) = k$.

6.6 The Case: $1 < \gcd(p-1, n+1) < n+1$. For convenience, set $g := \gcd(p-1, n+1)$. We investigate here the inequality in Proposition 6.3. Note that since $n+1$ does not divide $p-1$, neither $(p-1)\omega_1$ nor $(p-1)\omega_n$ lie in the root lattice. So we restrict attention to ω_j with $1 < j < n$. As such, there is nothing to consider unless $n \geq 3$. Without a loss of generality assume that $j \leq (n+1)/2$. Furthermore, from Remark 6.3, we may assume that $j(p-1)$ is divisible by $(n+1)$.

Consider the function $f(j) = \left[2 \left(\frac{p-1}{n+1} \right) - 1 \right] j(n+1-j)$, which is a quadratic polynomial in the variable j . For our purposes, we want to minimize $f(j)$. This evidently occurs when j is minimal (for $j \leq (n+1)/2$). So we consider the case that j is minimal such that $j(p-1)$ is divisible by $(n+1)$. This implies that, $n+1 = gj$, where $g = \gcd(n+1, p-1)$. With this substitution, using (6.3.5), the inequality in Proposition 6.3 may be rewritten as

$$(6.6.1) \quad i \geq [2(p-1) - (n+1)]j \left(1 - \frac{1}{g} \right).$$

Suppose first that $n + 1 = gj$ is odd (and $n \geq 4$). Then, both g and j must be odd. Therefore, $g \geq 3$ and $3 \leq j \leq (n + 1)/2$. Hence,

$$j \left(1 - \frac{1}{g}\right) \geq 3 \left(1 - \frac{1}{3}\right) = 2.$$

Equation (6.6.1) allows us to conclude that

$$\begin{aligned} i &\geq 2[2(p - 1) - (n + 1)] \\ &= 4(p - 1) - 2(n + 1) \\ &= 2(p - 1) + 2(p - 1) - 2(n + 1) \\ &= 2(p - 1) + 2(p - n - 2) \\ &\geq 2(p - 1) = 2p - 2 \end{aligned}$$

since $p \geq n + 2$. So we get a bound on i which is strictly larger than $2p - 3$.

Consider now the case that $n + 1 = gj$ is even (and $n \geq 3$). Since $p - 1$ is even, g is necessarily even. In particular, $g \geq 2$. Since $g \neq n + 1$, we also have $j \geq 2$. Suppose first that $j \geq 4$. Then

$$j \left(1 - \frac{1}{g}\right) \geq 4 \left(1 - \frac{1}{2}\right) = 2.$$

And so the same argument as in the $n + 1$ -odd case would show that $i \geq 2p - 2$.

Suppose next that $j = 3$ and $g > 2$. Then $g \geq 4$. Then,

$$j \left(1 - \frac{1}{g}\right) \geq 3 \left(1 - \frac{1}{4}\right) = \frac{9}{4} > 2.$$

We conclude that $i > 2p - 2$.

Suppose next that $j = 3$ and $g = 2$. Then $n + 1 = 6$ and $p - 1 > n + 1 = 6$. So $p \geq 11$ (as p is prime). Here we get

$$\begin{aligned} i &\geq [2(p - 1) - (n + 1)]j \left(1 - \frac{1}{g}\right) \\ &= [2(p - 1) - 6]3 \left(1 - \frac{1}{2}\right) \\ &= 3[(p - 1) - 3] \\ &= 3p - 12 \\ &= 2p - 3 + p - 9 \\ &\geq 2p - 1 \end{aligned}$$

since $p \geq 11$.

Suppose next that $j = 2$, i.e., $n + 1 = 2g$. Since $p - 1 > n + 1$ and g divides $p - 1$, we must have $p - 1 \geq 3g$. Write $p - 1 = (3 + m)g$ for an integer $m \geq 0$. Here we get

$$\begin{aligned}
i &\geq [2(p - 1) - (n + 1)]j \left(1 - \frac{1}{g}\right) \\
&= [2(p - 1) - 2g]2 \left(1 - \frac{1}{g}\right) \\
&= 4(p - 1) - \frac{4(p - 1)}{g} - 4g + 4 \\
&= 2(p - 1) + 2(p - 1) - \frac{4(p - 1)}{g} - 4g + 4 \\
&= 2(p - 1) + 2(3 + m)g - 4(3 + m) - 4g + 4 \\
&= 2(p - 1) + 2g - 8 + m(2g - 4) \\
&\geq 2(p - 1) + 2g - 8
\end{aligned}$$

since $m \geq 0$ and $g \geq 2$. If $g \geq 4$, then we conclude that $i \geq 2p - 2$.

However, if $g = 2$, we can only conclude that $i \geq 2p - 6$. This happens when $n + 1 = gj = 4$ or $n = 3$. Note that for $n = 3$, we either have $g = 2$ or $g = 4$ with the latter case falling into the $n + 1$ divides $p - 1$ category. The case of $n = 3$ will be dealt with specifically in Section 6.11. We summarize our findings in the following proposition.

Proposition . *Suppose Φ is of type A_n with $n \geq 4$. Suppose further that $p > n + 2$ and $1 < \gcd(p - 1, n + 1) < n + 1$. Let $\lambda = p\omega_j + w \cdot 0 \in X(T)_+$ for $2 \leq j \leq n - 1$ and $w \in W$. Then $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $0 < i \leq 2p - 3$.*

6.7 The Case: $p - 1 = n + 1$. Under this condition, we can explicitly construct a weight λ with $H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$. Let $\lambda = p\omega_1 + w \cdot 0$ where $w = s_1 s_2 s_3 \dots s_n$ with s_i denoting the simple reflection corresponding to the i th simple root. Then $-w \cdot 0$ is the sum of all n positive roots containing α_1 . In other words, $-w \cdot 0 = (n + 1)\omega_1$. So $\lambda = p\omega_1 + w \cdot 0 = p\omega_1 - (n + 1)\omega_1 = \omega_1$. Since $\ell(w) = n = p - 2$, by Proposition 3.2,

$$\dim H^{p-2}(G, H^0(\omega_1) \otimes H^0(\omega_1^*)^{(1)}) = \sum_{u \in W} (-1)^{\ell(u)} P_0(u \cdot \omega_1 - \omega_1) = 1.$$

Note that $\omega_1^* = \omega_n$ and one can similarly argue that $H^{p-2}(G, H^0(\omega_n) \otimes H^0(\omega_n^*)^{(1)}) = k$.

Theorem . *Suppose Φ is of type A_n with $n \geq 2$ and $p - 1 = n + 1$. Then*

- (a) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < p - 2$;
- (b) $H^{p-2}(G(\mathbb{F}_p), k) = k \oplus k$.

Proof. Part (a) follows from Theorem 4.4. For part (b), from the discussion above, we know that $H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = k$ for $\lambda = \omega_1$ or $\lambda = \omega_n$. We claim that if $H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$ for a dominant weight λ , then $\lambda = \omega_1$ or ω_n . To see this, write $\lambda = p\mu + w \cdot 0$ for some $\mu \in X(T)_+$ and $w \in W$. By Corollary 3.5(b), μ must be a fundamental dominant weight. Apply Proposition 6.3 with $p - 1 = n + 1$. The proposition

gives that, for $1 < j < n$,

$$i \geq j(n+1-j) = j(p-1-j) \geq 2(p-1-2) = 2p-6 = (p-2) + (p-4).$$

For $n \geq 3$, $p = n+2 \geq 5$ and so this gives $i > p-2$. Hence, $\mu = \omega_1$ or ω_n . In other words $\lambda = p\omega_1 + w \cdot 0$ or $\lambda = p\omega_n + w \cdot 0$, respectively. From the proof of Corollary 3.5, we observe that in order to have $H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$, when $-w \cdot 0$ is expressed as a sum of distinct positive roots, one of those roots must be $\tilde{\alpha} = \alpha_1 + \alpha_2 + \cdots + \alpha_n$. From Observation 2.2(B), it follows that $\ell(w) \geq n = p-2$. Applying Proposition 3.2, we see that

$$\dim H^{p-2}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = \sum_{u \in W} (-1)^{\ell(u)} P_0(u \cdot \lambda - \mu).$$

One has non-zero cohomology only if $u \cdot \lambda - \mu = 0$ which can only happen if $\lambda = \mu$, which gives the claim.

Since the only dominant weight less than ω_1 or ω_n is the zero weight, $H^i(G, k) = 0$ for $i > 0$, and ω_1 and ω_n lie in different linkage classes, the discussion in Section 2.8 gives part (b). \square

6.8 The Case: $\gcd(p-1, n+1) = n+1 < p-1$. The case $p-1 = n+1$ is excluded since that was dealt with in Section 6.7. Since $n+1$ divides $p-1$, $(p-1)\omega_j$ lies in the root lattice for all $1 \leq j \leq n$, and we need to allow $\lambda = p\omega_j + w \cdot 0$ for all j in our general argument.

Write $p-1 = d(n+1)$ for an integer $d \geq 2$. We can rewrite the inequality in Proposition 6.3 (see also (6.3.5)) as

$$\begin{aligned} i &\geq 2j(p-1) - 2j^2 \left(\frac{p-1}{n+1} \right) - j(n+1) + j^2 \\ &= 2(p-1) + (2j-2)(p-1) - 2j^2 \left(\frac{p-1}{n+1} \right) - j(n+1) + j^2 \\ &= 2(p-1) + [(2j-2)(n+1) - 2j^2] \left(\frac{p-1}{n+1} \right) - j(n+1) + j^2 \\ &= 2(p-1) + [(2j-2)(n+1) - 2j^2]d - j(n+1) + j^2. \end{aligned}$$

For $j = 1$ (or $j = n$), this inequality allows for a value of $i < 2p-3$. This will be discussed more in the next section. For this section, we focus on the case $2 \leq j \leq n-1$. By default, we need $n \geq 3$. For such j , the least value of the right hand side above occurs when $j = 2$ (or $j = n-1$). Substituting $j = 2$, the above inequality becomes

$$(6.8.1) \quad i \geq 2(p-1) + (2n-6)d - 2n + 2.$$

Since $d \geq 2$, (6.8.1) becomes

$$i \geq 2(p-1) + 2n - 10.$$

If $n \geq 5$, then $i \geq 2p-2$. If $n = 4$ with $d \geq 3$, then (6.8.1) becomes

$$i \geq 2(p-1) + (2*4-6)3 - 2*4 + 2 = 2p-2.$$

If $n = 4$ and $d = 2$, then $p-1 = 2(4+1) = 10$ or $p = 11$ and we can only say that $i \geq 2p-6$. This case will be considered in Section 6.12. For $n = 3$, notice that the value of

d is irrelevant in (6.8.1). Irrespective of d , we conclude that $i \geq 2p - 6$. This case will be discussed in Section 6.11. We summarize the conclusions of this section in the following.

Proposition . *Suppose Φ is of type A_n with $n \geq 4$. Suppose further that $p > n + 2$ and $\gcd(p-1, n+1) = n+1$. If $n = 4$, assume further that $p \neq 11$. Let $\lambda = p\omega_j + w \cdot 0 \in X(T)_+$ for $2 \leq j \leq n-1$ and $w \in W$. Then $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $0 < i \leq 2p - 3$.*

6.9 The Case: $\gcd(p-1, n+1) = n+1 < p-1$, **continued.** In this section we investigate the case of $\lambda = p\omega_1 + w \cdot 0$ (or symmetrically, $\lambda = p\omega_n + w \cdot 0$).

Lemma . *Suppose that Φ is of type A_n . Let $\lambda = p\omega_1 + w \cdot 0 \in X(T)_+$ or $\lambda = p\omega_n + w \cdot 0 \in X(T)_+$ and $\mu = p\omega_j + v \cdot 0 \in X(T)_+$ for $1 \leq j \leq n$ and $v, w \in W$. Then $H^i(G, H^0(\mu) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $0 < i < (p-1-n)n - (n+1-j)j + \ell(v)$.*

Proof. We give the argument for ω_1 . An analogous argument works for ω_n . If $\lambda = p\omega_1 + w \cdot 0$ is dominant, then a direct computation shows that $w = s_k s_{k-1} \dots s_1$ and $\lambda = (p-k-1)\omega_1 + \omega_{k+1}$, where $0 \leq k \leq n$. Here we are using the conventions $s_0 = 1$ and $\omega_{n+1} = 0$. Moreover, we have the following equations for the formal characters

$$\text{char}(V((p-k)\omega_1) \otimes V(\omega_k)) = \text{char } V((p-k)\omega_1 + \omega_k) + \text{char } V((p-k-1)\omega_1 + \omega_{k+1}).$$

As discussed earlier, the module $\text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j)$ has a good filtration. We conclude that

$$\begin{aligned} & \dim \text{Hom}_G(V(p\omega_1 + w \cdot 0), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j)) \\ & \leq \dim \text{Hom}_G(V((p-k)\omega_1) \otimes V(\omega_k), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j)) \\ & = \dim \text{Hom}_G(V((p-k)\omega_1), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j) \otimes H^0(\omega_{n+1-k})), \end{aligned}$$

where $k = \ell(w)$.

Note that $\langle \omega_j - u(\omega_k), \alpha_l^\vee \rangle \geq -1$ for all $u \in W$ and $1 \leq l \leq n$. It follows from [KLT] that $R^j \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes (\omega_j + u(\omega_{n+1-k}))) = 0$ for $j > 0$. Using the long exact sequence that one obtains from a B -filtration of $H^0(\omega_{n+1-k})$ one concludes that

$$\begin{aligned} & \dim \text{Hom}_G(V(p\omega_1 + w \cdot 0), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j)) \\ & \leq \dim \text{Hom}_G(V((p-k)\omega_1), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j) \otimes H^0(\omega_{n+1-k})) \\ & = \frac{1}{|\text{Stab}_W(\omega_k)|} \sum_{u \in W} \dim \text{Hom}_G(V((p-k)\omega_1), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j - u(\omega_k))). \end{aligned}$$

The Weyl module $V((p-k)\omega_1)$ has one-dimensional weight spaces [J, II 2.16]. A theorem of Kostant [Hum1, 24.2] implies that

$$\sum_{m \geq 0} \sum_{x \in W} (-1)^{\ell(x)} P_m(x \cdot ((p-k)\omega_1) - \omega_j + u(\omega_k)) = 1.$$

According to [AJ, 3.8] and [KLT] we know that

$$\begin{aligned} & \sum_{x \in W} (-1)^{\ell(x)} P_m(x \cdot ((p-k)\omega_1) - \omega_j + u(\omega_k)) \\ & = \dim \text{Hom}_G(V((p-k)\omega_1), \text{ind}_B^G(S^m(\mathbf{u}^*) \otimes \omega_j - u(\omega_k))) \geq 0, \end{aligned}$$

for all $m \geq 0$. Clearly, for $x \neq 1$,

$$\text{height}((p-k)\omega_1 - \omega_j + u(\omega_k)) > \text{height}(x \cdot ((p-k)\omega_1) - \omega_j + u(\omega_k)).$$

We conclude that

$$\sum_{x \in W} (-1)^{\ell(x)} P_m(x \cdot ((p-k)\omega_1) - \omega_j + u(\omega_k)) = \begin{cases} 1 & \text{if } m = \text{height}((p-k)\omega_1 - \omega_j + u(\omega_k)) \\ 0 & \text{else.} \end{cases}$$

We have

$$\begin{aligned} \text{height}((p-k)\omega_1 - \omega_j + u(\omega_k)) &\geq \text{height}((p-k)\omega_1 - \omega_j - \omega_{n+1-k}) \\ &= \frac{(p-k)n}{2} - \frac{(n+1-j)j}{2} - \frac{(n+1-k)k}{2}. \end{aligned}$$

Hence,

$$\dim \text{Hom}_G(V(p\omega_1 + w \cdot 0), \text{ind}_B^G(S^m \mathbf{u}^* \otimes \omega_j)) = 0,$$

if

$$m < \frac{(p-k)n - (n+1-j)j - (n+1-k)k}{2}$$

where $k = \ell(w)$.

Setting $m = \frac{i-\ell(v)}{2}$ and applying Lemma 3.1 and (3.2.1) yields

$$H^i(G, H^0(p\omega_1 + w \cdot 0) \otimes H^0((p\omega_j + v \cdot 0)^*)^{(1)}) = 0$$

for

$$\frac{i-\ell(v)}{2} < \frac{(p-k)n - (n+1-j)j - (n+1-k)k}{2}.$$

Hence, one obtains vanishing for

$$(6.9.1) \quad i < (p-k)n - (n+1-k)k - (n+1-j)j + \ell(v).$$

Since the global minimum of $(p-k)n - (n+1-k)k$ on the closed interval $[1, n]$ occurs at $k = n$, we get the claimed vanishing for

$$i \leq (p-1-n)n - (n+1-j)j + \ell(v).$$

□

Proposition . *Suppose Φ is of type A_n with $n \geq 3$. Suppose further that $p > n + 2$ and $\gcd(p-1, n+1) = n+1$. Let $\lambda = p\omega_1 + w \cdot 0 \in X(T)_+$ or $\lambda = p\omega_n + w \cdot 0$ with $w \in W$. Then $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $0 < i \leq 2p-3$.*

Proof. Equation (6.9.1) implies, for the case $\lambda = \mu$, vanishing for $0 < i < (p-k)n - (n+1-k)k - n + \ell(v) = (p-k)n - (n-k)k - n$. Again the global minimum occurs at $k = n$ and one obtains vanishing for $0 < i \leq (p-1-n)n$. Observe that $p \geq 2(n+1) + 1 = 2n+3$. Now

$$\begin{aligned} (p-1-n)n &= 2p-3 + p(n-2) - n(n+1) + 3 \\ &\geq 2p-3 + (2n+3)(n-2) - n^2 - n + 3 \\ &= 2p-3 + (n-1)^2 - 4 \\ &\geq 2p-3 \end{aligned}$$

with equality if and only if $n = 3$ and $p = 2n + 3$. This case does not occur. □

6.10 The Case: $n = 2$. Assume for this subsection that Φ is of type A_2 with $p > 3$. From Proposition 6.1 we know that $H^{2p-3}(G(\mathbb{F}_p), k) \neq 0$. Proposition 6.5 implies that $2p - 3$ is indeed the lowest bound unless 3 divides $p - 1$. Note that the case $p - 1 = 3$ is not possible for a prime p . If 3 divides $p - 1$, then the only possible non-zero cohomology in lower degrees would come from weights of the form $\lambda = p\omega_1 + w \cdot 0$ or the dual case $\lambda = p\omega_2 + w \cdot 0$. It follows from Lemma 6.9 (see also the proof of Proposition 6.9) that $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $i < 2p - 6$. Moreover, using the arguments of Lemma 6.9 one can show that $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $i \leq 2p - 6$ unless $\lambda = (p - 3)\omega_j$. In that case $\sum_{u \in W} (-1)^{\ell(u)} P_{p-4}(u \cdot ((p - 3)\omega_j) - \omega_j) = 1$ because the height of $(p - 4)\omega_j$ is exactly $p - 4$. Proposition 3.2 now says that $H^{2p-6}(G, H^0((p - 3)\omega_j) \otimes H^0((p - 3)\omega_{(3-j)})^{(1)}) = k$, $j = 1, 2$. Note that ω_1 and ω_2 are in different linkage classes. We conclude the following from Theorem 2.8(A) and the linkage discussion in Section 2.8.

Proposition . *Suppose Φ is of type A_2 and $p > 3$.*

- (a) *If 3 divides $p - 1$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 6$;
 - (ii) $H^{2p-6}(G(\mathbb{F}_p), k) = k \oplus k$.
- (b) *If 3 does not divide $p - 1$, then $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 3$.*

6.11 The case $n = 3$. Let Φ be of type A_3 with $p > 4$. The case $p = 5$ is included in Proposition 6.7. For the remainder of this section we assume that $p > 5$.

Lemma . *Suppose that Φ is of type A_3 with $p > 4$. Then*

$$\sum_{u \in W} (-1)^{\ell(u)} P_{p-5}(u \cdot ((p - 4)\omega_2) - \omega_2) = 1.$$

Proof. Observe that $2\omega_2 = \alpha_1 + 2\alpha_2 + \alpha_3 = \epsilon_1 + \epsilon_2 - \epsilon_3 - \epsilon_4$. For $u \cdot ((p - 4)\omega_2) - \omega_2$ to be a sum of positive roots one needs $u(2\omega_2)$ to be a sum of positive roots. This is the case if and only if either $u(2\omega_2) = 2\omega_2$ or $u(2\omega_2) = \alpha_1 + \alpha_3$. But we can rule out the second case because here $u \cdot ((p - 4)\omega_2) - \omega_2 = ((p - 5)/2)(\alpha_1 + \alpha_3) - \alpha_2 + u \cdot 0$. This is clearly not the sum of positive roots. It follows that we only have to consider $u \in \text{Stab}_W(\omega_2) = \{1, s_1, s_3, s_1 s_3\}$ and that $u \cdot ((p - 4)\omega_2) - \omega_2 = (p - 5)\omega_2 + u \cdot 0$. A direct computation now shows that

$$P_{p-5}((p - 5)\omega_2 + u \cdot 0) = \begin{cases} (p - 5)/2 & \text{if } u = 1 \\ (p - 7)/2 & \text{if } u = s_1, s_3, s_1 s_3. \end{cases}$$

Hence, $\sum_{u \in W} (-1)^{\ell(u)} P_{p-5}(u \cdot ((p - 4)\omega_2) - \omega_2) = \sum_{u \in \text{Stab}_W(\omega_2)} (-1)^{\ell(u)} P_{p-5}(u \cdot ((p - 4)\omega_2) - \omega_2) = 1$. \square

By Proposition 6.1 we know that $H^{2p-3}(G(\mathbb{F}_p), k) \neq 0$. Note that $\gcd(p - 1, n + 1) = \gcd(p - 1, 4) = 2$ or 4 . If $\gcd(p - 1, 4) = 2$, then by Remark 6.3, the only possible non-zero cohomology in lower degrees would come from weights of the form $\lambda = p\omega_2 + w \cdot 0$. On the other hand, if $\gcd(p - 1, 4) = 4$, then Proposition 6.9 gives the same conclusion.

Suppose $\lambda = p\omega_2 + w \cdot 0$. It follows from Proposition 6.3 that $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $i < 2p - 6$ and that $H^{2p-6}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ unless $\lambda = (p - 4)\omega_2$. That occurs

if $w \cdot 0 = -4\omega_2$. By direct computation one finds that $\ell(w) = 4$. By Proposition 3.2 and Lemma 6.11 above, we have

$$\begin{aligned} \dim H^{2p-6}(G, H^0((p-4)\omega_2) \otimes H^0((p-4)\omega_2^*)^{(1)}) \\ &= \sum_{u \in W} (-1)^{\ell(u)} P_{\frac{(2p-6)-4}{2}}(u \cdot ((p-4)\omega_2) - \omega_2) \\ &= \sum_{u \in W} (-1)^{\ell(u)} P_{p-5}(u \cdot ((p-4)\omega_2) - \omega_2) = 1. \end{aligned}$$

Combining this with Theorem 2.8(A) one obtains that $H^{2p-6}(G(\mathbb{F}_p), k) = k$. We summarize our findings for $\Phi = A_3$ below.

Proposition . *Suppose Φ is of type A_3 and $p > 4$.*

- (a) *If $p = 5$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < p - 2$;
 - (ii) $H^{p-2}(G(\mathbb{F}_p), k) = k \oplus k$.
- (b) *If $p > 5$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 6$;
 - (ii) $H^{2p-6}(G(\mathbb{F}_p), k) = k$.

6.12 The case $n = 4$ and $p = 11$. Assume for this section that Φ is of type A_4 with $p = 11$. Then $\gcd(p-1, n+1) = \gcd(10, 5) = 5$ and $2p-3 = 19$. It follows from Proposition 6.9 and Proposition 6.4 that the only non-zero cohomology in degrees lower than 19 has to come from weights of the form $\lambda = 11\omega_2 + w \cdot 0 = 6\omega_2$ or $\lambda = 11\omega_3 + w \cdot 0 = 6\omega_3$. In these cases we could potentially achieve non-vanishing in degree 18. The weights ω_2 and ω_3 are dual. We give the argument for ω_2 . An analogous argument works for ω_3 . Here $\ell(w) = 6$. According to Proposition 3.2, to have $H^{18}(G, H^0((p-5)\omega_2) \otimes H^0((p-5)\omega_3)^{(1)}) \neq 0$ one needs $\sum_{u \in W} (-1)^{\ell(u)} P_6(u \cdot (6\omega_2) - \omega_2) \neq 0$. The lemma below (Lemma 6.12) rules this case out.

According to Proposition 6.1, $H^{19}(G(\mathbb{F}_p), k) \neq 0$. Again it follows from Proposition 6.9 and Proposition 6.4 that any cohomology in degree 19 other than the one coming from Proposition 6.1 has to come from weights of the form $\lambda = 11\omega_2 + w \cdot 0$ or $\lambda = 11\omega_3 + w \cdot 0$. Suppose $\lambda = 11\omega_2 + w \cdot 0$. Again, the ω_3 case is analogous. Proposition 3.2 implies that cohomology in odd degrees has to come from weights with corresponding $w \in W$ of odd length. Now equation (6.3.1) shows that this can only happen if $\ell(w) = 5$. Note that here $M_2(-w \cdot 0) = \ell(w) = 5$. The only such weight is $\lambda = 6\omega_2 + \tilde{\alpha}$. By Proposition 3.2 it suffices to show that $\sum_{u \in W} (-1)^{\ell(u)} P_7(u \cdot (6\omega_2 + \tilde{\alpha}) - \omega_2) = 0$. We conclude from the lemma below that $H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ for $0 < i < 2p - 3$ and further that $H^{2p-3}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) = 0$ unless $\lambda = (p-n)\tilde{\alpha} = 7\tilde{\alpha}$.

Lemma . *Suppose that Φ is of type A_4 with $p = 11$. Then*

- (a) $\sum_{u \in W} (-1)^{\ell(u)} P_6(u \cdot (6\omega_2) - \omega_2) = 0$,
- (b) $\sum_{u \in W} (-1)^{\ell(u)} P_7(u \cdot (6\omega_2 + \tilde{\alpha}) - \omega_2) = 0$.

Proof. (a) Observe that $5\omega_2 = 3\alpha_1 + 6\alpha_2 + 4\alpha_3 + 2\alpha_4 = 3\epsilon_1 + 3\epsilon_2 - 2\epsilon_3 - 2\epsilon_4 - 2\epsilon_5$. For $u \cdot (6\omega_2) - \omega_2$ to be a sum of positive roots one needs $u(5\omega_2)$ to be a sum of positive roots.

This is the case if and only if either $u(5\omega_2) = 5\omega_2$ or $u(5\omega_2) = 3\alpha_1 + \alpha_2 + 4\alpha_3 + 2\alpha_4$. Let us consider the second case. Here u is an element of the co-set $s_2 \cdot \text{Stab}_W(\omega_2)$. Note that for any $u \in s_2 \cdot \text{Stab}_W(\omega_2)$ the expression $-u \cdot 0$ contains some positive multiple of the root α_2 . Therefore $u \cdot (6\omega_2) - \omega_2 = 3\alpha_1 + 4\alpha_3 + 2\alpha_4 + u \cdot 0$ is not a sum of positive roots. It suffices therefore to look at $\sum_{u \in \text{Stab}_W(\omega_2)} (-1)^{\ell(u)} P_6(5\omega_2 + u \cdot 0)$. A straightforward but tedious calculation now shows that $\sum_{u \in \text{Stab}_W(\omega_2)} (-1)^{\ell(u)} P_6(5\omega_2 + u \cdot 0) = 0$.

(b) As in part (a) for $u \cdot (6\omega_2 + \tilde{\alpha}) - \omega_2$ to be the sum of positive roots one needs either $u \in \text{Stab}_W(\omega_2)$ or $u \in s_2 \cdot \text{Stab}_W(\omega_2)$. However, the second case results in weights that cannot be written as sums of 7 positive roots. As in part (a) it suffices therefore to look at $\sum_{u \in \text{Stab}_W(\omega_2)} (-1)^{\ell(u)} P_7(6\omega_2 + u \cdot \alpha_0)$ which can be shown to be zero.

Both parts of the lemma can also be readily checked by using computer software such as MAGMA [BC, BCP]. \square

6.13 Summary for $r = 1$. The following theorem addresses (1.1.1) and (1.1.2) for type A_n when $r = 1$.

Theorem . *Suppose Φ is of type A_n with $n \geq 2$. Suppose further that $p > n + 1$.*

- (a) *(Generic case) If $p > n + 2$ and $n > 3$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 3$;
 - (ii) $H^{2p-3}(G(\mathbb{F}_p), k) = k$.
- (b) *If $p = n + 2$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < p - 2$;
 - (ii) $H^{p-2}(G(\mathbb{F}_p), k) = k \oplus k$.
- (c) *If $n = 2$ and 3 divides $p - 1$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 6$;
 - (ii) $H^{2p-6}(G(\mathbb{F}_p), k) = k \oplus k$.
- (d) *If $n = 2$ and 3 does not divide $p - 1$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 3$;
 - (ii) $H^{2p-3}(G(\mathbb{F}_p), k) = k$.
- (e) *If $n = 3$ and $p > 5$, then*
 - (i) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < 2p - 6$;
 - (ii) $H^{2p-6}(G(\mathbb{F}_p), k) = k$.

Proof. Let $\lambda = p\mu + w \cdot 0 \in X(T)_+$ for $\mu \in X(T)_+$ and $w \in W$. Based on the discussion in Section 2, our goal has been to determine the least $i > 0$ such that

$$H^i(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0.$$

According to Proposition 6.1(a), we know that the weight $\lambda = p\tilde{\alpha} - n\tilde{\alpha}$ gives a non-zero cohomology class in degree $2p - 3$.

By Proposition 3.4(b), if $\langle \mu, \tilde{\alpha}^\vee \rangle \geq 2$, then $i \geq 2p - 3$. Hence, the only way to obtain a smaller i is for μ to be a fundamental weight. That case has been dealt with in previous sections, from which parts (a)(i), (b), (c), (d)(i), and (e) follow. It remains to show parts (a)(ii) and (d)(ii). From Proposition 6.4, $\lambda = p\tilde{\alpha} - n\tilde{\alpha} = p(\omega_1 + \omega_n)$ is the only weight with $H^{2p-3}(G, H^0(\lambda) \otimes H^0(\lambda^*)^{(1)}) \neq 0$. The result follows by Proposition 6.1(a) and Theorem 2.8(A). \square

6.14 Generic result for $r > 1$. The following theorem addresses (1.1.1) and (1.1.2) for type A_n when $r > 1$.

In summarizing the results for the $r = 1$ situation in Section 6.13 we encountered many special cases for small ranks and small primes. One would expect even more non-generic behavior for $r > 1$. For example, it is easily seen that in the case $p = n + 2$ non-vanishing already occurs in degree $r(p - 2)$, i.e., $H^{r(p-2)}(G(\mathbb{F}_p), k) \neq 0$. To give a complete answer many case-by-case arguments will be necessary, most of them rather lengthy and intricate. For brevity we limit ourselves here to the generic case.

Theorem . *Suppose Φ is of type A_n with $n \geq 4$ and $p > 2(n + 1)$. Suppose further that $p > 11$ whenever $n = 4$. Then*

- (a) $H^i(G(\mathbb{F}_p), k) = 0$ for $0 < i < r(2p - 3)$;
- (b) $H^{r(2p-3)}(G(\mathbb{F}_p), k) \neq 0$.

Proof. Assume that $\text{Ext}_G^i(V(\lambda)^{(r)}, H^0(\lambda)) \neq 0$. By Proposition 4.2 there exists a sequence of non-zero weights $\lambda = \gamma_0, \gamma_1, \dots, \gamma_{r-1}, \gamma_r = \lambda \in X(T)_+$ and nonnegative integers l_1, l_2, \dots, l_r such that

- (i) $i = \sum_{j=1}^r l_j$,
- (ii) $\text{Ext}_G^{l_j}(V(\gamma_j)^{(1)}, H^0(\gamma_{j-1})) \neq 0$, for $1 \leq j \leq r$ and
- (iii) $\gamma_j = p\delta_j + w_j \cdot 0$ for some $u_j \in W$ and nonzero $\delta_j \in X(T)_+$.

Next we apply our discussion in Section 6.3 to the pair of weights γ_j, γ_{j-1} and obtain from (6.3.2)

$$(6.14.1) \quad l_j \geq 2pM_{m(\delta_j)}(\delta_j) + 2M_{m(\delta_j)}(w_j \cdot 0) - 2M_{m(\delta_{j-1})}(\delta_{j-1}) - M_{m(\delta_{j-1})}(w_{j-1} \cdot 0).$$

It follows from $\delta_r = \delta_0$ and $w_r = w_0$ that

$$\begin{aligned} i = \sum_{j=1}^r l_j &\geq \sum_{j=1}^r 2pM_{m(\delta_j)}(\delta_j) + 2M_{m(\delta_j)}(w_j \cdot 0) - 2M_{m(\delta_{j-1})}(\delta_{j-1}) - M_{m(\delta_{j-1})}(w_{j-1} \cdot 0) \\ &= \sum_{j=1}^r 2(p-1)M_{m(\delta_j)}(\delta_j) + M_{m(\delta_j)}(w_j \cdot 0) \\ &\geq \sum_{j=1}^r 2(p-1)M_{m(\delta_j)}(\delta_j) + M_{m(\delta_j)}(-2\rho), \end{aligned}$$

where the last inequality follows from Observation 6.3(f). We define

$$(6.14.2) \quad d_j := l_j + 2M_{m(\delta_{j-1})}(\delta_{j-1}) - 2M_{m(\delta_j)}(\delta_j) - M_{m(\delta_j)}(w_j \cdot 0) + M_{m(\delta_{j-1})}(w_{j-1} \cdot 0).$$

Then $i = \sum_{j=1}^r d_j$ and $d_j \geq 2(p-1)M_{m(\delta_j)}(\delta_j) + M_{m(\delta_j)}(-2\rho)$.

According to Observation 6.3(e), we know that $M_{m(\delta_j)}(-2\rho) = -N_{m(\delta_j)}$. Moreover, using the arguments in the proofs of Propositions 6.2, 6.3 and 6.4 one obtains the following bounds:

$$(6.14.3) \quad d_j \geq \begin{cases} \left[2 \left(\frac{p-1}{n+1} \right) - 1 \right] (n+1 - m(\delta_j)) m(\delta_j) & \text{if } \delta_j = \omega_{m(\delta_j)}, \\ 2p-3 & \text{if } \delta_j = \omega_1 + \omega_n, \\ 2(p-1) + 2(n-1) \left[\frac{p-1}{n+1} - 1 \right] & \text{else.} \end{cases}$$

Note that, as observed in Section 6.4, the last expression is larger than $2p - 3$ under the assumptions on p and n .

If $m(\delta_j) \neq 1, n$, the expression $\left[2 \left(\frac{p-1}{n+1}\right) - 1\right](n+1 - m(\delta_j))m(\delta_j)$ is least when $m(\delta_j) = 2$ or $n - 1$ (see Section 6.8). Hence,

$$\begin{aligned} d_j &\geq 2(p-1) + (2n-6) \left(\frac{p-1}{n+1}\right) - 2n + 2 \\ &\geq 2(p-1) + 2(2n-6) - 2n + 2 \\ &= 2(p-1) + 2n - 10. \end{aligned}$$

If $n \geq 5$ one obtains $d_j \geq 2(p-1)$. For $n = 4$ one can show that $d_j \geq 2p - 3$ whenever $p > 11$. We conclude that under our assumptions $d_j \geq 2p - 3$ unless $\delta_j = \omega_1$ or ω_n .

Next assume that $\delta_j = \omega_1$ or ω_n . The above methods produce the lower bound

$$(6.14.4) \quad d_j \geq 2(p-1) - 2 \left(\frac{p-1}{n+1}\right) - n,$$

which is not sufficient. Note that ω_1 or ω_n are not in the root lattice. Therefore $\delta_{j-1} \neq \omega_1 + \omega_n = \tilde{\alpha}$. We will distinguish two cases.

Case 1: δ_{j-1} is not a fundamental weight.

In this case (6.14.3) yields

$$(6.14.5) \quad d_{j-1} \geq 2(p-1) + 2(n-1) \left[\frac{p-1}{n+1} - 1\right].$$

Adding (6.14.4) and (6.14.5) produces

$$\begin{aligned} d_j + d_{j-1} &\geq 2(2p-3) + 2 + 2(n-2) \left(\frac{p-1}{n+1}\right) - 3n + 2 \geq 2(2p-3) + 4(n-2) - 3n + 4 \\ &= 2(2p-3) + n - 4 \geq 2(2p-3). \end{aligned}$$

Case 2: $\delta_{j-1} = \omega_s$, $1 \leq s \leq n$.

We apply Lemma 6.9 to obtain from equation (6.9.1)

$$l_j \geq (p-k)n - (n+1-k)k - (n+1-s)s + \ell(w_{j-1}),$$

where $k = \ell(w_j) = -M_1(w_j \cdot 0)$. By (6.14.2)

$$d_j \geq l_j + 2M_s(\omega_s) - 2M_1(\omega_1) - M_1(w_j \cdot 0) + M_s(w_{j-1} \cdot 0).$$

From Observation 2.2(B) (see Section 6.3), we know that $\ell(w_{j-1}) \geq -M_s(w_{j-1} \cdot 0)$ and clearly $M_s(\omega_s) \geq M_1(\omega_1)$. This yields

$$d_j \geq (p-k)n - (n+1-k)k + k - (n+1-s)s = (p-k)n - (n-k)k - (n+1-s)s.$$

As a function of k and s the above obtains its minimum at $k = n$ and $s = (n+1)/2$. Hence

$$d_j \geq (p-n)n - \frac{(n+1)^2}{4} = 2p - 3 + (n-2)p - n^2 - \frac{(n+1)^2}{4} + 3.$$

Using $p \geq 2n + 3$ one obtains

$$\begin{aligned} d_j &\geq 2p - 3 + (n - 2)(2n + 3) - n^2 - \frac{n^2 + 2n + 1}{4} + 3 \\ &= 2p - 3 + n^2 - 2n + 1 - \frac{n^2 - 2n + 1}{4} - 4 = 2p - 3 + \frac{3(n - 1)^2}{4} - 4. \end{aligned}$$

For $n \geq 4$ this yields $d_j > 2p - 3$.

From above, we see that $d_j \geq 2p - 3$ unless $\delta_j = \omega_1$ or ω_n and δ_{j-1} is not a fundamental weight. If $d_j \geq 2p - 3$ for each $1 \leq j \leq r$, we have

$$i \geq \sum_{j=1}^r d_j \geq r(2p - 3)$$

and part (a) follows. Suppose now that $d_j < 2p - 3$ for some j . Let t be the largest such j and suppose that $t > 1$. Then

$$i \geq \sum_{j=1}^r d_j = \sum_{t+1}^r d_j + d_t + d_{t-1} + \sum_{j=1}^{t-2} d_j \geq (r - t)(2p - 3) + 2(2p - 3) + \sum_{j=1}^{t-2} d_j.$$

Consider the remaining sum $\sum_{j=1}^{t-2} d_j$, again identify the largest j with $d_j < 2p - 3$, and repeat this decomposition. Continuing in this manner the claim follows except possibly if we reach a point when the largest j with $d_j < 2p - 3$ is $j = 1$. So we are done if $d_1 \geq 2p - 3$.

Suppose now that $\delta_1 = \omega_1$ or ω_n and δ_0 is not a fundamental weight so that we could have $d_1 < 2p - 3$. Note that since δ_1 is a fundamental weight, $d_2 \geq 2p - 3$. Recall that $\delta_0 = \delta_r$. Therefore, we can use the Case 1 argument to show that $d_1 + d_r \geq 2(2p - 3)$. Further, since $d_2 \geq 2p - 3$, the above argument can successfully be used to show that $\sum_{j=2}^{r-1} d_j \geq (r - 2)(2p - 3)$. Hence,

$$i \geq d_1 + d_r + \sum_{j=2}^{r-1} d_j \geq 2(2p - 3) + (r - 2)(2p - 3) = r(2p - 3).$$

The proof of part (a) is now complete, and part (b) is derived from Proposition 6.2(b). \square

REFERENCES

- [AJ] H. H. Andersen, J. C. Jantzen, Cohomology of induced representations for algebraic groups, *Math. Ann.*, **269**, (1984), 487–525.
- [AM] A. Adem, R.J. Milgram, *Cohomology of Finite Groups*, Springer-Verlag, Berlin Heidelberg, 1994.
- [BNP1] C. P. Bendel, D. K. Nakano, C. Pillen, On comparing the cohomology of algebraic groups, finite Chevalley groups, and Frobenius kernels, *J. Pure & Appl. Algebra*, **163**, no. 2, (2001), 119-146.
- [BNP2] C. P. Bendel, D. K. Nakano, C. Pillen, Extensions for finite Chevalley groups I, *Adv. Math.*, **183**, (2004), 380–408.
- [BNP3] C. P. Bendel, D. K. Nakano, C. Pillen, Extensions for finite Chevalley groups II, *Trans. Amer. Math. Soc.*, **354**, no. 11, (2002), 4421–4454.
- [BNP4] C. P. Bendel, D. K. Nakano, C. Pillen, Extensions for Frobenius kernels, *J. Algebra*, **272**, (2004), 476-511.
- [BNP5] C. P. Bendel, D. K. Nakano, C. Pillen, Extensions for finite groups of Lie type: twisted groups, in *Finite Groups 2003*, eds. Ho et al., de Gruyter, New York, 2004, pp. 29-46.
- [BNP6] C. P. Bendel, D. K. Nakano, C. Pillen, Extensions for finite groups of Lie Type II: filtrating the truncated induction functor, *Cont. Math.*, **413**, (2006), 1-23.

- [BNP7] C. P. Bendel, D. K. Nakano, C. Pillen, Second cohomology groups for Frobenius kernels and related structures, *Adv. Math.*, **209**, (2007), 162-197.
- [BC] W. Bosma, J. Cannon, *Handbook on Magma Functions*, Sydney University, 1996.
- [BCP] W. Bosma, J. Cannon, C. Playhoust, The Magma Algebra System I: The User Language, *J. Symbolic Computation*, **3/4** no. 24, (1997), 235-265.
- [C] J. F. Carlson, The cohomology of irreducible modules over $SL(2, p^n)$, *Proc. London Math. Soc.* (3), **47**, (1983), 480-492.
- [F] E. M. Friedlander, Computations of K -theories of finite fields, *Topology*, **15**, (1976), 87-109.
- [FP] E. M. Friedlander, B. J. Parshall On the cohomology of algebraic and related finite groups, *Invent. Math.*, **74**, (1983), 85-118.
- [H] H. L. Hiller, Cohomology of Chevalley groups over finite fields, *J. Pure Appl. Alg.*, **16**, (1980), 259-263.
- [Hum1] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, 1972.
- [Hum2] J. E. Humphreys, *Modular Representations of Finite Groups of Lie Type*, Amer. Math. Soc., Providence, RI, 2003.
- [J] J. C. Jantzen, *Representations of Algebraic Groups*, American Mathematical Society, Mathematical Surveys and Monographs, **107**, 2003.
- [KLT] S. Kumar, N. Lauritzen, J. F. Thomsen, Frobenius splitting of cotangent bundles of flag varieties, *Invent. Math.*, **136**, (1999), 603-621.
- [Q] D. G. Quillen, On the cohomology and K -theory of finite fields, *Ann. Math.*, **96**, (1972), 551-586.
- [W] C. B. Wright, Second cohomology groups for algebraic groups and their Frobenius kernels, preprint, 2009.

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF WISCONSIN-STOUT, MENOMONIE, WI 54751, USA

E-mail address: bendelc@uwstout.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GA 30602, USA

E-mail address: nakano@math.uga.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SOUTH ALABAMA, MOBILE, AL 36688, USA

E-mail address: pillen@jaguar1.usouthal.edu