

Lehmer's Question, Knots and Surface Dynamics

BY DANIEL S. SILVER AND SUSAN G. WILLIAMS†

*Department of Mathematics and Statistics University of South Alabama,
Mobile, AL 36688, U.S.A.*

(Received 22 March 2006; revised 15 January 2007)

Abstract

Lehmer's question is equivalent to one about generalized growth rates of Lefschetz numbers of iterated pseudo-Anosov surface homeomorphisms. One need consider only homeomorphisms that arise as monodromies of fibered knots in lens spaces $L(n, 1)$, $n > 0$. Lehmer's question for Perron polynomials is equivalent to one about generalized growth rates of words under free group endomorphisms.



1. Introduction

The Mahler measure $M(f)$ of a nonzero polynomial $f(x) = c_n x^n + \cdots + c_1 x + c_0$ ($c_n \neq 0$) $\in \mathbb{C}[x]$ is

$$M(f) = |c_n| \prod_{j=1}^n \max(|r_j|, 1),$$

where r_1, \dots, r_n are the roots of f . Lehmer's question asks whether the Mahler measure of a polynomial with integer coefficients can be arbitrarily close but not equal to 1. D.H. Lehmer posed the question in [30]. He could do no better than 1.17628..., a value that he achieved with the remarkable polynomial $L(x) = x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1$. No value closer but not equal to 1 has yet been found despite extensive computer searches [5, 7, 37, 39]. History and status of various searches can be viewed at M. Mossinghoff's web page www.cecm.sfu.ca/~mjm/Lehmer.

Mahler measure can be defined, albeit in a different manner, for polynomials of more than one variable. Lehmer's question turns out to be equivalent to the identical question for polynomials in any fixed number of variables (see [6]).

Lehmer's polynomial evaluated at $x = -t$ is the Alexander polynomial of the $(-2, 3, 7)$ -pretzel knot, a fibered hyperbolic knot with striking properties [27]. Motivated by this, the authors applied techniques of algebraic dynamics [42] to provide a topological interpretation of the Mahler measure of Alexander polynomials of knots and links in terms of homology of finite abelian branched covering spaces [43]. Many hyperbolic knots and links with simple complements have Alexander polynomials with remarkably small Mahler measure [44].

All of the knots and links that arise in [44] are fibered, which means that their complements are mapping tori of surface homeomorphisms. It is natural to ask whether Lehmer's

† Both authors partially supported by NSF grant DMS-0304971.

question is equivalent to a question about surface homeomorphisms. We answer this affirmatively in Section 3. In Section 4 we go further and show that Lehmer’s question is equivalent to a question about generalized growth rates of Lefschetz numbers of iterated pseudo-Anosov homeomorphisms. The growth rates that we use are a generalization of the usual one, introduced by Kronecker and Hadamard.

Section 5 explores free group endomorphisms, growth rates of words and connections with Lehmer’s question. The final section examines connections with braids.

2. Growth rates of sequences

Let $\mathbf{a} = \{a_n\}_{n=1}^\infty$ be a sequence of complex numbers. For each positive integer k , we define the k th growth rate $\text{GR}^{(k)}(\mathbf{a})$ to be

$$\limsup_{n \rightarrow \infty} \left| \det \begin{pmatrix} a_n & a_{n+1} & \cdots & a_{n+k-1} \\ a_{n+1} & a_{n+2} & \cdots & a_{n+k} \\ \vdots & & & \vdots \\ a_{n+k-1} & a_{n+k} & \cdots & a_{n+2k-2} \end{pmatrix} \right|^{1/n}.$$

We define the 0th growth rate $\text{GR}^{(0)}(\mathbf{a})$ to be 1. Notice that $\text{GR}^{(1)}(\mathbf{a})$ is the usual growth rate

$$\limsup_{n \rightarrow \infty} |a_n|^{1/n},$$

which appears for example in [3]. The general growth rates $\text{GR}^{(k)}(\mathbf{a})$ were studied by Kronecker and Hadamard in the late 19th century. The determinants $H_{n,k}$ above are known as *Hankel determinants*. The following proposition is a consequence [18] (see also [9, p. 335]).

PROPOSITION 2.1. *The sequence \mathbf{a} is the sequence of coefficients of a rational power series*

$$\sum_{n=0}^\infty a_n t^n = R(t) / \prod_{i=1}^d (1 - \lambda_i t) \tag{2.1}$$

if and only if for some N the Hankel determinants $H_{n,k}$ vanish for all $k > d$ and $n \geq N$. In this case, $\text{GR}^{(k)}(\mathbf{a}) = 0$ if $k > d$. If no $1 - \lambda_i t$ divides $R(t)$ and $|\lambda_1| \geq \cdots \geq |\lambda_d|$, then $\text{GR}^{(1)}(\mathbf{a}) = |\lambda_1|$, $\text{GR}^{(2)}(\mathbf{a}) = |\lambda_1 \lambda_2|$, \dots , $\text{GR}^{(d)}(\mathbf{a}) = |\lambda_1 \lambda_2 \cdots \lambda_d|$. Consequently, $\max_k \text{GR}^{(k)}(\mathbf{a}) = M(f)$, where $f(t) = \prod_{i=1}^d (t - \lambda_i)$.

By a *linear recurrence* we will mean a homogenous linear recurrence relation with constant coefficients, that is, a relation

$$a_{n+d} + c_{d-1}a_{n+d-1} + \cdots + c_1a_{n+1} + c_0a_n = 0,$$

$n \geq 0$, with *characteristic polynomial*

$$f(t) = t^d + c_{d-1}t^{d-1} + \cdots + c_1t + c_0 = \prod_{i=1}^d (t - \lambda_i).$$

(An excellent reference on this topic is [11].) The sequence $\mathbf{a} = \{a_n\}$ satisfies this linear recurrence if and only if it is the sequence of coefficients of the rational power series (2.1), where $R(t) = b_{d-1}t^{d-1} + \cdots + b_1t + b_0$ with $b_k = \sum_{i=0}^k a_i c_{k-i}$. The *minimal polynomial* of the sequence \mathbf{a} is the unique minimal degree characteristic polynomial of a linear recurrence

for \mathbf{a} . Note that if the rational function (2.1) is in reduced form then the minimal polynomial of \mathbf{a} is $t^m \prod_{i=1}^d (t - \lambda_i)$ for some non-negative integer m . Thus we have:

PROPOSITION 2.2. *If $\mathbf{a} = \{a_n\}_{n=1}^\infty$ is a linearly recurrent sequence with minimal polynomial $f(t)$ then*

$$\max_k \text{GR}^{(k)}(\mathbf{a}) = M(f).$$

The following result will be needed in Section 4.

PROPOSITION 2.3. *Suppose $\mathbf{a} = \{a_n\}_{n=1}^\infty$ and $\mathbf{b} = \{b_n\}_{n=1}^\infty$ are linearly recurrent sequences with $|a_n - b_n|$ bounded. Then the minimal polynomials of \mathbf{a} and \mathbf{b} have the same roots outside the unit circle, with the same multiplicities. In particular, $\max_k \text{GR}^{(k)}(\mathbf{a}) = \max_k \text{GR}^{(k)}(\mathbf{b})$.*

Proof. Set $\mathbf{c} = \{c_n\} = \{a_n - b_n\}$. Then \mathbf{c} is also linearly recurrent. The power series $\sum_{n=0}^\infty c_n t^n$ converges on $|t| < 1$. Writing this power series as a rational function as in (2.1), we see that the roots of the minimal polynomial of \mathbf{c} must lie in $|t| \leq 1$. From the general theory of linear recurrences, we can write $a_n = \sum p_i(n)\alpha_i^n$, $b_n = \sum q_i(n)\beta_i^n$, $c_n = \sum r_i(n)\gamma_i^n$, where α_i , β_i and γ_i range over the distinct roots of the minimal polynomials of \mathbf{a} , \mathbf{b} and \mathbf{c} respectively, and p_i , q_i and r_i are polynomials of degree one less than the multiplicity of the corresponding root. Any set of functions on \mathbb{N} of the form $n^k \lambda^n$ with distinct $(k, \lambda) \in \mathbb{N} \times \mathbb{C}$ is linearly independent. Hence the terms in the sums $\sum p_i(n)\alpha_i^n$ and $\sum q_i(n)\beta_i^n$ corresponding to roots of modulus greater than one must be identical. The desired result is immediate.

Remark 2.4. If we assume in addition to the hypothesis of Proposition 2.3 that \mathbf{a} and \mathbf{b} are integer sequences, then the difference $a_n - b_n$ is eventually periodic (see [38, part VIII, problem 158]).

3. Fibered links

Let $\ell = \ell_1 \cup \dots \cup \ell_d \subset M^3$ be an oriented link in an oriented closed 3-manifold with exterior $X = M \setminus \text{int}N(\ell)$. Here $N(\ell) = \ell \times \mathbb{D}^2$ denotes a regular neighbourhood of ℓ . The link is *fibered* if the projection $\partial N(\ell) = \ell \times \mathbb{S}^1 \rightarrow \mathbb{S}^1$ extends to a locally trivial fibration of X . In this case, X is homeomorphic to a *mapping torus* $S \times [0, 1]/\{(s, 0) \sim (h(s), 1)\}$, for some compact orientable surface (*fiber*) S and homeomorphism (called a *monodromy*) $h: S \rightarrow S$, which is unique up to isotopy.

A monodromy h induces an automorphism h_* of $H_1(S; \mathbb{R}) \cong \mathbb{R}^d$. The characteristic polynomial $\text{char}(h_*) = \det(h_* - tI)$ is a monic reciprocal integral polynomial of even degree that is well defined up to multiplication by $\pm t^n$. When M is the 3-sphere, it can be obtained from the classical multivariable Alexander polynomial $\Delta_\ell(t_1, \dots, t_d)$ of ℓ by replacing each variable t_i by t , and dividing by $(t - 1)^{d-1}$ if $d > 1$. Details can be found for example in [24, proposition 7.3.10].

A surface automorphism $h: S \rightarrow S$ is *periodic* if h^r is isotopic to the identity for some $r > 0$. It is *reducible* if it is isotopic to an automorphism that leaves invariant some essential 1-manifold of S . A theorem of Thurston [48] asserts that if h is neither periodic nor reducible and S is hyperbolic, then after a suitable isotopy, h leaves invariant a pair of transverse singular measured foliations (\mathcal{F}^u, μ^u) , (\mathcal{F}^s, μ^s) such that

$$f(\mathcal{F}^u, \mu^u) = (\mathcal{F}^u, \lambda \mu^u), \quad f(\mathcal{F}^s, \mu^s) = (\mathcal{F}^s, \lambda^{-1} \mu^s),$$

for some $\lambda > 1$. Consequently, h expands leaves of \mathcal{F}^u by a factor of λ , and it contracts

leaves of \mathcal{F}^s by λ^{-1} . Such an automorphism is said to be *pseudo-Anosov*. Details for closed surfaces can be read in [2]. Modifications needed for surfaces with boundary can be found in [22] and [29]. Another result of Thurston [47] states that the mapping torus of a surface automorphism $h: S \rightarrow S$ is a hyperbolic 3-manifold if and only if h is isotopic to a pseudo-Anosov homeomorphism.

The following lemma is proved in [2, section 5] for the case of a closed surface. The argument applies equally well to surfaces with connected boundary. For the reader's convenience, we repeat the argument with necessary modifications.

LEMMA 3.1. *Assume that $h: S \rightarrow S$ is a homeomorphism of an compact connected orientable surface, closed or with connected boundary. If $\text{char}(h_*)$ is not composite, cyclotomic or a polynomial in t^r for any $r > 1$, then h is neither periodic nor reducible.*

Proof. If h is periodic, then $h_*^n = 1$ for some $n > 0$, which implies that $\text{char}(h_*)$ is cyclotomic, a contradiction.

Assume that h is reducible. Then after isotopy, $h(C) = C$ for some essential 1-submanifold C . The components of C are disjoint simple closed curves, none homotopic to a point or ∂S , and no two homotopic to each other. We distinguish two cases:

(1) Some component C_1 is non-separating. Orient C_1 . It follows that $0 \neq [C_1] \in H_1(S)$. For some $n > 0$, $h_*^n[C_1] = [C_1]$. Hence some eigenvalue of h_* is a root of unity, and $\text{char}(h_*)$ is cyclotomic, a contradiction.

(2) Each component of C separates S . There is a component S_0 of $S \setminus C$ with frontier equal to a single component of C . If S has boundary, then we can assume that S_0 does not contain ∂S . Note that $H_1(S_0)$ must be nontrivial. Consider the least $r > 0$ such that $h^r(S_0) = S_0$. Then

$$H_1(S) = H_1(S_0) \oplus H_1(h(S_0)) \oplus \cdots \oplus H_1(h^{r-1}(S_0)) \oplus H_1(\hat{G}),$$

where \hat{G} , possibly empty, is obtained from $G = S \setminus (S_0 \cup h(S_0) \cup \cdots \cup h^{r-1}(S_0))$ by capping off boundary components other than ∂S .

Since the summands $H_1(h^i(S_0))$ are nontrivial, irreducibility of $\text{char}(h_*)$ implies that $H_1(\hat{G})$ is trivial. It follows that $r > 1$, for otherwise C consists of a single curve that is homotopic through G to the boundary of S or to a point if ∂S is empty. Since the automorphism h_* permutes the summands $H_1(h^i(S_0))$ cyclically, $\text{char}(h_*)$ has the form $\det(B - t^r I)$, for some matrix B . Hence $\text{char}(h_*)$ is a polynomial in t^r , a contradiction.

The Alexander polynomial $\Delta(k)(t)$ of a fibered knot $k \subset \mathbb{S}^3$ is an integral monic polynomial that satisfies the conditions: (1) $\Delta(k)(t)$ is reciprocal; (2) $\Delta(k)(t)$ has even degree; (3) $\Delta(k)(1) = \pm 1$. Conversely, any such polynomial arises from a fibered knot. When investigating Lehmer's question, it suffices to consider integral monic polynomials satisfying conditions (1) and (2) (see [45]). Unfortunately, it is not known if (3) can also be assumed. In order to circumvent this problem, Hironaka suggested that one consider reduced Alexander polynomials of fibered links.

By [23], for any integral monic polynomial $f(t)$ satisfying conditions (1) and (2), there exists a fibered 2-component link ℓ with $\Delta(\ell)(t, t) \doteq f(t)$. (Here, and throughout, ' \doteq ' indicates equality up to multiplication by monomials $\pm t^n$.) The polynomial $\Delta(\ell)(t, t)$ is sometimes called the *reduced Alexander polynomial* of ℓ , denoted by $\Delta_{\text{red}}(\ell)(t)$.

Example 3.2. Recall that $L(t)$ denotes Lehmer's polynomial (see Introduction). It is well known that $L(-t)$ is the Alexander polynomial of the $(-2, 3, 7)$ -pretzel knot, a fibered hyperbolic knot. Addition of a meridional circle produces a fibered 2-component link ℓ with




Fig. 1. Fibered link from Kanenobu's construction.

$\Delta_{\text{red}}(\ell)(t) = L(-t)$. Kanenobu's general construction produces a much more complicated link ℓ with $\Delta_{\text{red}}(\ell)(t) = L(-t)$. The general form of Kanenobu's links appears in Figure 1. The link is displayed as the boundary of a fiber surface S . A knot with Alexander polynomial $L(-t)$ results from the choice $n = 5$, $p_1 = -11$, $p_2 = 43$, $p_3 = -69$, $p_4 = 36$ and $p_5 = 1$.

In general, the links that Kanenobu constructs have the feature that at least one component is unknotted (shown in Figure 1).

The definition of Alexander polynomial for knots in \mathbb{S}^3 extends for knots k in rational homology 3-spheres M^3 [50]. As in the classical case, when k is fibered (that is, $M \setminus k$ fibers over the circle), the Alexander polynomial $\Delta(k)(t)$ coincides with $\text{char}(h_*)$, where h is a monodromy. It is well defined up to multiplication by $\pm t^i$.

Let p, q be coprime integers. We recall that the lens space $L(p, q)$ is the union of two solid tori V_1, V_2 identified along their respective boundary tori T_1, T_2 by a homeomorphism from V_2 to V_1 that sends the homology class of μ_2 to that of $p\lambda_1 + q\mu_1$. Here (μ_i, λ_i) are respective meridian-longitude bases of T_i , $i = 1, 2$. It is a standard fact that $\pi_1(L(p, q)) \cong \mathbb{Z}/p$. Also, $L(1, 1)$ is the 3-sphere \mathbb{S}^3 . Additional background can be found, for example, in [40].

THEOREM 3.3. *Lehmer's question is equivalent to the following. For any $\epsilon > 0$, does there exist a fibered hyperbolic knot k in a lens space $L(n, 1)$, $n > 0$, such that $1 < M(\Delta(k)(t)) < 1 + \epsilon$?*

Proof. We need consider only polynomials $f(t)$ which are not composite, cyclotomic or a polynomial in t^r for any $r > 1$. Kanenobu's construction produces a fibered 2-component link $\ell = \ell_1 \cup \ell_2$ with monodromy $h: S \rightarrow S$ such that $\text{char}(h_*) = (t - 1)f(t)$. Capping off the boundary component of S corresponding to an unknotted component, say ℓ_2 , produces a homeomorphism $\hat{h}: \hat{S} \rightarrow \hat{S}$ of a surface \hat{S} with a single boundary component, and $\text{char}(h_*) = f(t)$. By Lemma 3.1, \hat{h} is isotopic to a pseudo-Anosov homeomorphism.

We prove that the mapping torus of \hat{h} is the exterior of a knot k in a lens space $L(n, 1)$, where $n = f(1)$. (By [21], n is the linking number of the two components of ℓ .) Let C_1, C_2 denote the boundary components of S . Each is a longitude of the corresponding component ℓ_1, ℓ_2 . Since ℓ_2 is trivial, we may assume after isotopy that it is the standard unknot; let $V = \ell_2 \times \mathbb{D}^2$ be a regular neighborhood of ℓ_2 . Let V_1 be the complementary solid torus in \mathbb{S}^3 . Take λ_1 to be a meridian of ℓ_2 , μ_1 a nontrivial curve in ∂V_1 that bounds a disk in V .

The curve ℓ_1 has winding number n in V_1 . Note that since $f(t)$ is irreducible and non-cyclotomic, n is nonzero. The mapping torus of \hat{h} is obtained from $V_1 \setminus \text{int}N(\ell_1)$ by attaching another solid torus $V_2 = \mathbb{D}^2 \times \mathbb{S}^1$ with $\partial \mathbb{D}^2 \times \theta = C_2 \times \theta$, for each $\theta \in \mathbb{S}^1$. It suffices to show that C_2 meets ∂V_1 in a curve homotopic to $\mu_1 + n\lambda_1$.

It is clear that C_2 wraps once in the μ_1 direction. The longitude C_1 of ℓ_1 wraps n times in the λ_1 direction. Since the fiber S is a homology between C_1 and C_2 , it follows immediately that C_2 also wraps n times. This completes the argument that ℓ_1 is a fibered knot in $L(n, 1)$.

Since $L(-n, 1)$ is homeomorphic to $L(n, 1)$, we may assume without loss of generality that $n > 0$.

Finally, we observe that since the monodromy \hat{h} is pseudo-Anosov, the complement of ℓ_1 is hyperbolic.

Remark 3.4.

- (i) The knot k can be obtained directly from ℓ by surgery on the unknotted component ℓ_2 .
- (ii) We can cap off the remaining boundary component of \hat{S} and extend \hat{h} to a homeomorphism \bar{h} of a closed surface \bar{S} . The characteristic polynomials of \hat{h}_* and \bar{h}_* are equal. Lemma 3.1 implies that Lehmer's question is equivalent to a question about pseudo-Anosov homeomorphisms of closed orientable surfaces.

4. Dynamics

The periodic orbit structure of a surface homeomorphism contains much information about the dynamics of the map. The fixed points of a pseudo-Anosov homeomorphism h are isolated and hence can be counted. Since any iterate of h is also pseudo-Anosov, the cardinality of the fixed point set $\text{Fix}(h^n)$ is well defined for any integer n , and we denote it by F_n . One can regard $\text{Fix}(h^n)$ as the set of points of period n under h .

The *Lefschetz number* $L(h)$ of a pseudo-Anosov homeomorphism h is the sum of the indices of its fixed points $p \in \text{Fix}(h)$. (Lefschetz numbers are defined for more general maps.) If p is an isolated fixed point in the interior of S , then its *index* is the degree of $\text{id} - h : (S, S \setminus \{p\}) \rightarrow (S, S \setminus \{p\})$. When the fixed point is contained on the boundary of S , then one computes the index by first embedding S in Euclidean space as a neighbourhood retract and then extending h . Details can be found in [10] (see also [15]).

By [13], $2\chi(S) - 1 \leq \text{index}(p) \leq 1$. Moreover, the index at any non-prong singularity is ± 1 , and the number of all fixed points with index less than -1 is bounded below by $4\chi(S)$ [25].

We abbreviate the Lefschetz number $L(h^n)$ by L_n . In what follows, we consider the sequence $\mathbf{L} = \{L_n\}$.

Lefschetz numbers are homotopy invariants of h , and the well-known Lefschetz fixed point theorem implies that L_n is equal to the alternating sum of homology traces:

$$\begin{aligned} L_n &= \sum_{i=0}^2 (-1)^i \text{tr}[h_*^n : H_i(S; \mathbb{R}) \rightarrow H_i(S; \mathbb{R})] \\ &= \begin{cases} 1 - \text{tr} A^n, & \text{if } \partial S \neq \emptyset; \\ 2 - \text{tr} A^n, & \text{if } \partial S = \emptyset, \end{cases} \end{aligned}$$

where A is a matrix that represents $h_* : H_1(S, \mathbb{R}) \rightarrow H_1(S, \mathbb{R})$.

Let $f(t) = \det(h_* - tI)$. Then $\text{tr} A^n = \lambda_1^n + \dots + \lambda_d^n$, where $\lambda_1, \dots, \lambda_d$ are the roots (with possible repetition) of $f(t)$. The sequence $\mathbf{L} = \{L_n\}_{n=1}^\infty$ is linearly recurrent with minimal polynomial $p(t) = (t-1) \prod (t - \lambda_i)$, where the product ranges over distinct non-zero λ_i . By Proposition 2.2 we have:

PROPOSITION 4.1. *If h is a pseudo-Anosov homeomorphism and $f(t) = \det(h_* - tI)$, then*

$$\max_k \text{GR}^{(k)}(\mathbf{L}) \leq M(f),$$

with equality if f has no repeated roots.

Combining this with the proof of Theorem 3.3 gives:

COROLLARY 4.2. *Lehmer's question is equivalent to the following. For any $\epsilon > 0$, does there exist a pseudo-Anosov homeomorphism $h: S \rightarrow S$ such that $\max_k \text{GR}^{(k)}(\mathbf{L})$ is contained in the interval $(1, 1 + \epsilon)$?*

One can assume that h is a monodromy for a fibered 2-component link in \mathbb{S}^3 or a hyperbolic fibered knot in $L(n, 1)$, $n > 0$. Alternatively, Remark 3.4 implies that we can restrict our attention to homeomorphisms of closed orientable surfaces.

There exist pseudo-Anosov homeomorphisms h such that h_* is the identity map [49], and consequently all growth rates $\text{GR}^{(k)}(\mathbf{L})$ are trivial. For such homeomorphisms, nontrivial growth rates are achieved by replacing Lefschetz numbers by the number F_n of periodic points. It is a consequence of [33] and [14] that $\mathbf{F} = \{F_n\}_{n=1}^\infty$ is linearly recurrent. We sketch the proof for the convenience of the reader.

LEMMA 4.3. *If $h: S \rightarrow S$ is a pseudo-Anosov homeomorphism, then $\mathbf{F} = \{F_n\}$ satisfies a linear recurrence.*

Proof. By [14], the homeomorphism $h: S \rightarrow S$ admits a Markov partition. Consequently, there exists an $N \times N$ matrix

$$A = (a_{i,j})_{(i,j) \in N \times N} \tag{4.1}$$

with entries equal to 0 or 1 and a surjective map $p: \Omega \rightarrow S$, where

$$\Omega = \{(x_n)_{n \in \mathbb{Z}} \mid a_{x_n, x_{n+1}} = 1, n \in \mathbb{Z}\}$$

such that $p \circ \sigma = h \circ p$, where σ is the shift (to the left) of the sequence (x_n) of symbols. The homeomorphism σ is the *shift of finite type* with adjacency matrix A . For any n , the number of fixed points of σ^n is equal to the trace of A^n (see [4], for example). However, p need not be injective, and the trace of A^n can be larger than the number F_n of period n points of h . A combinatorial argument of Manning [33] can be used to correct the count. (For a homological approach, see [17].) One constructs additional shifts of finite type σ_i , $i = 1, \dots, m$, with adjacency matrices A_i and signs $\epsilon_i \in \{-1, 1\}$ such that for each n ,

$$F_n = \sum_{i=0}^m \epsilon_i \cdot \text{tr} A_i^n. \tag{4.2}$$

Let $p(t) = \prod (t - \lambda_j)$ where λ_j ranges over the distinct non-zero roots of the characteristic polynomials of A_0, \dots, A_m . Then F_n is linearly recurrent with minimal polynomial $p(t)$.

The invariant foliations are *orientable* if all of the intersections of any transverse arc with any leaf are in the same direction. For surfaces with connected boundary, the condition is equivalent to the requirement that the degree of every interior prong singularity is even.

The pseudo-Anosov monodromy of the $(-2, 3, 7)$ -pretzel knot has orientable invariant foliations (see remark following [31, theorem 9.7]). For it, $\max_k \text{GR}^{(k)}(\mathbf{F}) = \text{GR}^{(1)}(\mathbf{F})$ is equal to the Mahler measure of Lehmer's polynomial. More generally, we have:

THEOREM 4.4. Assume that $h: S \rightarrow S$ is a pseudo-Anosov homeomorphism with oriented invariant foliations. Then for any positive integer k ,

$$\text{GR}^{(k)}(\mathbf{F}) = \text{GR}^{(k)}(\mathbf{L}).$$

Combined with Proposition 4.1 this gives:

COROLLARY 4.5. Assume that $h: S \rightarrow S$ is a pseudo-Anosov homeomorphism with oriented invariant foliations and $f(t) = \det(h_* - tI)$. Then

$$\max_k \text{GR}^{(k)}(\mathbf{F}) \leq M(f),$$

with equality if f has no repeated roots.

Proof of Theorem 4.4. For any $n \geq 1$, all non-prong fixed points of h^n have the same index, $+1$ or -1 . One way to see this is to choose such a fixed point p , and orient the leaf L_p^u of \mathcal{F}^u containing p . The index at p is -1 if h preserves the orientation of L_p^u ; otherwise it is $+1$ (see [16, proposition 5.7]). Let q be any other non-prong fixed point of h^n , and let L_q^u be the leaf of \mathcal{F}^u containing it. Since L_p^u is dense in S , it passes arbitrarily close to q , always in the same direction, since \mathcal{F}^u is oriented. By continuity, the orientations of L_p^u and L_q^u are both preserved or else both reversed. Thus the index of q agrees with that of p .

If h reverses orientations of unstable leaves, then replace each F_n by $(-1)^{n-1}F_n$. Otherwise, replace each F_n by $-F_n$. In either case, the generalized growth rates are unchanged. Any remaining difference between F_n and L_n is due to prong singularities and hence is bounded. Proposition 2.3 completes the proof.

Remark 4.6. Theorem 3.3 of [41] states that for any pseudo-Anosov homeomorphism h with orientable foliations, the eigenvalues of h_* are the same as those of the matrix (4.1) including multiplicity, with the possible exception of some zeros and roots of unity. Corollary 4.5 above asserts that in the case that h has oriented invariant foliations, the generalized growth rates of the sequence $\{\text{trace } A^n\}$ are equal to those of \mathbf{F} .

5. Free group endomorphisms

Let G be any group with a finite set of generators g_1, \dots, g_m . The length $|g|$ of any $1 \neq g \in G$ is the minimum length of a word in $g_1^{\pm 1}, \dots, g_m^{\pm 1}$ expressing g . If ϕ is any endomorphism of G , then the growth rate $\text{GR}(\phi)$ is

$$\text{GR}(\phi) = \max_{1 \leq i \leq m} \limsup_{n \rightarrow \infty} |\phi^n(g_i)|^{1/n}.$$

It is independent of the generator set [3].

When $f: S \rightarrow S$ is a pseudo-Anosov homeomorphism, the growth rate $\text{GR}(f_{\mathbb{Z}})$ of the induced automorphism $f_{\mathbb{Z}}$ of $G = \pi_1(S)$ contains a wealth of information about the dynamics of f . For example, $\text{GR}(f_{\mathbb{Z}})$ coincides with the first growth rate $\text{GR}^{(1)}(\mathbf{F})$, and it is an upper bound for the modulus of every eigenvalue of $f_*: H_1(S; \mathbb{R}) \rightarrow H_1(S; \mathbb{R})$ (see [14] and [3]). Hence

$$\text{GR}(f_{\mathbb{Z}}) = \text{GR}^{(1)}(\mathbf{F}) \geq \text{GR}^{(1)}(\mathbf{L}). \quad (5.1)$$

Topological entropy $h_{\text{top}}(f)$ is a measure of complexity defined for any continuous map $f: X \rightarrow X$ of a compact metric space. The entropy of any periodic map is zero, while that of a pseudo-Anosov map is strictly positive. Although topological entropy is difficult to

compute in general, when X is a surface and f is pseudo-Anosov, $h_{\text{top}}(f)$ can be computed as $\log \text{GR}(f_{\pm})$ [14].

Definition 5.1. Let G be a group generated by g_1, \dots, g_m , and let $\phi: G \rightarrow G$ be an endomorphism. The k th generalized growth rate of ϕ (with respect to the given generating set) is

$$\text{GR}^{(k)}(\phi) = \max_{1 \leq i \leq m} \text{GR}^{(k)}(|\phi^n(g_i)|).$$

Example 5.2. In general these growth rates may depend on the generating set when $k > 1$. Consider the free group F of rank two generated by x, y and the endomorphism $\phi: F \rightarrow F$ sending x to x^3 and y to y^2 . It is easy to see that $|\phi^n(x)| = 3^n$, $|\phi^n(y)| = 2^n$ and hence $\text{GR}^{(1)}(\phi) = 3$ while $\text{GR}^{(k)}(\phi) = 0$ for $k > 1$.

On the other hand, if we replace x, y with generators z, y , where $z = xy$, then $\phi^n(z) = x^{3^n} y^{2^n} = (zy^{-1})^{3^n-1} zy^{2^n-1}$ and $|\phi^n(z)| = 2 \cdot 3^n + 2^n - 2$. The latter is linearly recurrent with minimal polynomial $(t-1)(t-2)(t-3)$ and hence $\text{GR}^{(1)}(\phi) = 3$, $\text{GR}^{(2)}(\phi) = \text{GR}^{(3)}(\phi) = 6$ while $\text{GR}^{(k)}(\phi) = 0$ for $k > 3$.

Remark 5.3. If we define $\text{GR}^{(k)}(\phi)$ instead to be $\text{GR}^{(k)}(\sum_{i=1}^m |\phi^n(g_i)|)$ then the two generating sets in the above example give the same maximum growth rate. We do not know if this holds in general. It is easily seen that this definition reduces to the usual growth rate $\text{GR}(\phi)$ when $k = 1$.

An important case of Lehmer's question can be viewed as a question about free group endomorphisms.

A square matrix A with nonnegative real entries is *primitive* if there is an $N > 0$ such that all the entries of A^N are positive. The characteristic polynomial of any primitive integer matrix is a monic integral polynomial such that one root is a positive real number that is strictly larger than the modulus of the remaining roots. Such a polynomial is said to be a *Perron polynomial*, and the dominant root is called a *Perron number*. For background about Perron–Frobenius theory, the reader might consult [36].

It follows from equation (4.2) that the minimal polynomial $p(t)$ of \mathbf{F} is a product of Perron polynomials.

THEOREM 5.4. *Given any irreducible Perron polynomial $p(t)$ there exists an endomorphism ϕ of a finitely generated free group such that for any generator g_i , the sequence of word lengths $|\phi^n(g_i)|$ is linearly recurrent with minimal polynomial equal to $p(t)$ times a product of cyclotomic polynomials. Thus the maximum growth rate of the sequence is $M(p)$.*

Proof. For $\Lambda = (\lambda_1, \lambda_2, \dots, \lambda_d)$ a d -tuple of complex numbers, we define $\text{tr}(\Lambda^n) = \sum_{i=1}^d \lambda_i^n$ and $\text{tr}_n(\Lambda) = \sum_{k|n} \mu(n/k) \text{tr}(\Lambda^k)$ where μ is the Moebius function. (The second quantity is called the n th net trace of Λ .) The main theorem of [26] states that there exist a primitive integer matrix A and a non-negative integer ℓ such that $\text{char}(A) = t^\ell \prod_{i=1}^d (t - \lambda_i)$ if and only if:

- (i) the polynomial $\prod_{i=1}^d (t - \lambda_i)$ has integer coefficients;
- (ii) some λ_i is real and $\lambda_i > |\lambda_j|$ for $i \neq j$;
- (iii) $\text{tr}_n(\Lambda) \geq 0$ for all $n \geq 1$.

Let $p(t) = \prod_{i=1}^d (t - \lambda_i)$ be a Perron polynomial. Since one root dominates the moduli of the other roots, the net trace condition is satisfied for sufficiently large n . We observe that the k -tuple of k th roots of unity has n th net trace equal to k for $n = k$ and 0 otherwise.

Thus we can multiply $p(t)$ by a suitable product $\Phi(t)$ of cyclotomic polynomials to obtain a polynomial $p(t)\Phi(t)$ for which the tuple of roots satisfies the net trace condition for all $n \geq 1$. Then using [26], we can find a primitive integer $m \times m$ matrix $A = (a_{i,j})$ with characteristic polynomial equal to $t^\ell p(t)\Phi(t)$ for some ℓ .

Let F_m be the free group generated by g_1, \dots, g_m . Define $\phi: F_m \rightarrow F_m$ to be the endomorphism mapping g_i to $g_1^{a_{i,1}} g_2^{a_{i,2}} \cdots g_m^{a_{i,m}}$.

Only nonnegative exponents occur in any reduced word representing $\phi^n(g_i)$, for any $n \geq 1$ and $i = 1, \dots, m$. Hence the number of occurrences of g_j in $\phi^n(g_i)$ is equal to $(A^n)_{i,j}$. The Cayley-Hamilton theorem implies that for fixed i and j , the sequence $(A^n)_{i,j}$ satisfies a linear recurrence with characteristic polynomial $\text{char}(A)$. Hence for any i , so does the sequence of word lengths $|\phi^n(g_i)|$, which is equal to $\sum_j (A^n)_{i,j}$. Since A is primitive, the sequence grows exponentially at a rate given by the dominant eigenvalue. Hence the minimal polynomial of the sequence is equal to the Perron polynomial $p(t)$ times some factor of $\Phi(t)$.

If $\ell = 0$, then ϕ is injective since its image abelianizes to a free abelian group of rank m . In such a case, we cannot expect $\phi: F \rightarrow F$ in Theorem 5.3 to be induced by an automorphism of an orientable surface, since in such case the set of roots of $p(t)$ must be closed under inversion (see [46]). Instead we ask:

Question 5.5. Can $\phi: F \rightarrow F$ in Theorem 5.4 be chosen to be a free group automorphism?

It is well known that given any matrix $A \in GL_m(\mathbb{Z})$, there exists a free group automorphism $\phi \in \text{Aut}(F_m)$ that abelianizes to A (see [32]). We say an endomorphism of the free group is *positive* if there is a generating set for which the image of every generator is a positive word, that is, a product of positive powers of generators.

Question 5.6. Suppose that $A \in GL_m(\mathbb{Z})$ has nonnegative entries. Does there exist a positive automorphism of the free group F_m that abelianizes to A ?

We conclude the section with a partial answer to Question 5.6.

PROPOSITION 5.7. *Suppose that $A \in GL_2(\mathbb{Z})$ has nonnegative entries. Then there is a positive automorphism of the free group F of rank 2 that abelianizes to A .*

Proof. Write

$$A = \begin{pmatrix} p_0 & p_1 \\ q_0 & q_1 \end{pmatrix}.$$

We may assume (exchanging the two generators if necessary) that either $p_0 - p_1$ or $q_0 - q_1$ is positive; the other is necessarily nonnegative.

Inductively, for $k \geq 0$ we write

$$\begin{pmatrix} p_k \\ q_k \end{pmatrix} = d_{k+1} \begin{pmatrix} p_{k+1} \\ q_{k+1} \end{pmatrix} + \begin{pmatrix} p_{k+2} \\ q_{k+2} \end{pmatrix},$$

where d_{k+1} is chosen to be the largest integer with p_{k+2}, q_{k+2} both nonnegative. We stop when $n = k + 2$ satisfies $p_n = 0$ or $q_n = 0$. We see easily that each matrix

$$\begin{pmatrix} p_k & p_{k+1} \\ q_k & q_{k+1} \end{pmatrix}$$

is in $GL_2(\mathbb{Z})$ with $p_k - p_{k+1}$ or $q_k - q_{k+1}$ positive. In particular, if $p_n = 0$ then $q_n = p_{n-1} = 1$, while if $q_n = 0$ then $p_n = q_{n-1} = 1$.

Let F_2 be the free group of rank 2 generated by a, b . Set $u_0 = a^{p_n} b^{q_n}$, $u_1 = a^{p_{n-1}} b^{q_{n-1}}$ (these generate F_2), and $u_{k+1} = u_k^{d_{n-k}} u_{k-1}$ for $k = 1, \dots, n - 1$. Define $\phi: F_2 \rightarrow F_2$ by $\phi(a) = u_n$, $\phi(b) = u_{n-1}$. We see by descent that u_k is in the image of ϕ for all k , and so ϕ is onto. Any surjective endomorphism of F_2 is an automorphism.

6. Braids and Lehmer’s question

An n -braid can be regarded as an orientation-preserving homeomorphism of the punctured disk $D_n = \mathbb{D}^2 \setminus \{n \text{ points}\}$, two n -braids regarded as the same if they are isotopic rel boundary. Alternatively, one can regard an n -braid as an isotopy class of a map of the pair $(\mathbb{D}^2, \{n \text{ points}\})$. The latter point of view enables us to recover the usual geometric picture of an n -braid by tracing the image of the set of n points as an isotopy rel boundary is performed from the map of \mathbb{D}^2 to the identity.

The collection of all n -braids forms a group B_n under concatenation. As usual, we let σ_i denote the final map of an isotopy of D_n that exchanges the i th and $i + 1$ st puncture, dragging the i th puncture around the $i + 1$ st in the clockwise direction. Then B_n is generated by $\sigma_1, \dots, \sigma_{n-1}$. The center of B_n is cyclic, generated by a full twist $\Delta_n^2 = (\sigma_{n-1} \sigma_{n-2} \cdots \sigma_1)^n$.

The Thurston–Nielsen classification of surface homeomorphisms applies to n -braids. One calls $\beta \in B_n$ periodic, reducible or pseudo-Anosov if it can be represented by a homeomorphism of D_n with that property.

Given an n -braid β , its *braid entropy* h_β is the infimum $\inf\{h_{\text{top}}(f) \mid f \text{ represents } \beta\}$. Braid entropy is a conjugacy invariant. Moreover, the braid entropy of $\beta \in B_n$ is equal to that of $\Delta_n^{2k} \beta$, for any integer k . If $f: D_n \rightarrow D_n$ is any representative of β , then h_β can be computed as $\log \text{GR}(f_\#)$.

The reduced Burau representation $\beta \mapsto \tilde{B}_\beta$ maps B_n to $\text{GL}(n - 1, \mathbb{Z}[t^{\pm 1}])$ via

$$\tilde{B}_{\sigma_1} = \begin{pmatrix} -t & 0 \\ 1 & 1 \\ & & I \end{pmatrix}, \quad \tilde{B}_{\sigma_i} = \begin{pmatrix} I & & & \\ & 1 & t & 0 \\ & 0 & -t & 0 \\ & 0 & 1 & 1 \\ & & & & I \end{pmatrix}, \quad \tilde{B}_{\sigma_{n-1}} = \begin{pmatrix} I & & \\ & -t & 0 \\ & 1 & 1 \end{pmatrix},$$

where I denotes an identity matrix of the appropriate size.

The closure $\hat{\beta}$ of an n -braid is the oriented link obtained by joining the top and bottom of each strand of β without introducing additional crossings. The strands of β are oriented coherently, from top to bottom. The following is well known (see [1]).

PROPOSITION 6.1. *The matrix $\tilde{B}_\beta - I$ is a Jacobian of the link $\hat{\beta}$. Moreover, $\det(\tilde{B}_\beta - I)$ is equal to $(1 + t + \cdots + t^{n-1}) \Delta_{\text{red}}(\hat{\beta})(t)$, where Δ_{red} is the reduced Alexander polynomial.*

As a consequence of results of Section 3 we have:

PROPOSITION 6.2. *Lehmer’s question is equivalent to the following. For any $\epsilon > 0$, does there exist a hyperbolic n -braid β such that $1 < M(\det(\tilde{B}_\beta - I)) < 1 + \epsilon$?*

The example below suggests that the relationship between Lehmer’s question and surface automorphisms is significant. We see that for $n = 3, 4$ and 5 , there exists a minimal entropy pseudo-Anosov n -braid $f: D_n \rightarrow D_n$ that closes to the $(-2, 3, 7)$ -pretzel knot.

In each case, the mapping torus of f is a hyperbolic 3-manifold M with $H_1 M \cong \mathbb{Z}^2$. There are many possible homomorphisms $\pi_1 M \rightarrow \mathbb{Z}$ with finitely generated kernel, homomorphisms that one might call fibering directions. (Fibering directions can be viewed as points of

fibered faces of a polyhedral unit ball of the Thurston norm. See [35].) One such direction yields the punctured disk D_n as fiber and f as monodromy. Other fibering directions result in pseudo-Anosov monodromies of possibly higher genus surfaces. In each of the three cases below, there exists a fibering direction for which the corresponding monodromy, capped off along the boundary of D_n , is that of the $(-2, 3, 7)$ -pretzel knot.

Example 6.3.

- (i) The braid $\sigma_1\sigma_2^{-1}$ is minimal among all pseudo-Anosov 3-braids in the forcing partial order [34, 20]. Hence it attains the minimum braid entropy ($= \log 2.61803 \dots$) among all pseudo-Anosov 3-braids. We note here that the closure of $\sigma_1\sigma_2^{-1}\Delta_3^4$, which has the same braid entropy, is the $(-2, 3, 7)$ -pretzel knot.
- (ii) The braid $\sigma_3\sigma_2\sigma_1^{-1}$ attains the minimum braid entropy ($= \log 2.29663 \dots$) among all pseudo-Anosov 4-braids [19]. The closure of $\sigma_3\sigma_2\sigma_1^{-1}\Delta_4^2$ is the $(-2, 3, 7)$ -pretzel knot.
- (iii) The braid $\sigma_1\sigma_2\sigma_3\sigma_4\sigma_1\sigma_2$ attains the minimum braid entropy ($= \log 1.72208 \dots$) among all pseudo-Anosov 5-braids [19]. The closure of $\sigma_1\sigma_2\sigma_3\sigma_4\sigma_1\sigma_2\Delta_5^2$ is again the $(-2, 3, 7)$ -pretzel knot.

We expect the mapping torus of a pseudo-Anosov homeomorphisms with small entropy to be a hyperbolic 3-manifold that is small in the sense of [8]; that is, composed of relatively few ideal tetrahedra. The examples above do not disappoint. The first and third have 4 tetrahedra, while the second has 5. Their volumes are $4.0597 \dots$, $3.6638 \dots$ and $4.8511 \dots$, respectively. Calculations were done with SnapPea. We note that the volume of the third example is the same as that of the Whitehead link complement, and is the smallest known volume of any hyperbolic 2-component link.

Acknowledgements. We are grateful to Vasilii Prokhorov for informing us about Hadamard's work. Mike Boyle, David Fried, Eriko Hironaka, Michael Kelly and Alexander Stoimenow also contributed helpful comments. We thank the referee for suggestions and corrections.

REFERENCES

- [1] J. S. BIRMAN. Braids, links and mapping class groups. *Ann. of Math. Stud.* **82** (1974).
- [2] S. A. BLEILER and A. J. CASSON. *Automorphisms of Surfaces after Nielsen and Thurston*. London Math. Soc. Student Texts **9** (Cambridge University Press, 1988).
- [3] R. BOWEN. *Entropy and the Fundamental Group* (Springer LNM **668** (1978), 21–29).
- [4] R. BOWEN and O. LANFORD. Zeta functions of restrictions of the shift transformation, in *Global Analysis. Proc. Symp. Pure Math.* **14** (1968).
- [5] D. W. BOYD. Reciprocal polynomials having small measure. *Math. Comp.* **35** (1980), 1361–1377.
- [6] D. W. BOYD. Speculations concerning the range of Mahler's measure. *Canad. Math. Bull.* **24** (1981), 453–469.
- [7] D. W. BOYD. Reciprocal polynomials having small Mahler measure II. *Math. Comp.* **53** (1989), 355–357.
- [8] P. J. CALLAHAN, J. C. DEAN and J. R. WEEKS. The simplest hyperbolic knots. *J. Knot Theory Ramifications* **8** (1999), 279–297.
- [9] P. DIENES. *The Taylor Series* (Dover Publications, 1957).
- [10] A. DOLD. *Lectures on Algebraic Topology, 2nd edition* (Springer Verlag, 1980).
- [11] G. EVEREST, A. VAN DER POORTEN, I. SHPARLINSKI and T. WARD. *Recurrence Sequences* (American Mathematical Society, 2003).
- [12] B. FARB, C. LEININGER and D. MARGALIT. In preparation.
- [13] A. FATHI and F. LAUDENBACH. Les feuilletages mesurés. *Astérisque* **66–67** (1979), 71–89.
- [14] A. FATHI and M. SHUB. Some dynamics of pseudo-Anosov diffeomorphisms. *Astérisque* **66–67** (1979), 181–207.

- [15] A. FEL'SHTYN. Dynamical zeta functions, Nielsen theory and Reidemeister torsion. *Mem. Amer. Math. Soc.* **699** (2000).
- [16] J. M. FRANKS. Homology and dynamical systems. *Conf. Board of the Math. Sciences Regional Conf. Series in Math.* **49** (Amer. Math. Soc., 1982).
- [17] D. FRIED. Finitely presented dynamical systems. *Ergodic Theory Dynam. Systems* **7** (1987), 489–507.
- [18] J. HADAMARD. Essai sur l'étude des fonctions données par leurs développements de Taylor. *J. de Mathématiques* (4) **8** (1892), 101–186.
- [19] J.-Y. HAM and W. T. SONG. The minimum dilatation of pseudo-Anosov 5-braids. Preprint, 2005, arXiv: math.GT/506295.
- [20] M. HANDEL. The forcing partial order on the three times punctured disk. *Ergodic Theory Dynam. Systems* **17** (1997), 593–610.
- [21] F. HOSOKAWA. On ∇ -polynomials of links. *Osaka Math. J.* **10** (1958), 273–282.
- [22] B. JIANG and J. GUO. Fixed points of surface diffeomorphisms. *Pacific J. Math.* **160** (1993), 67–89.
- [23] T. KANENOBU. Module d'Alexander des noeuds fibrés et polynôme de Hosokawa des lacements fibrés. *Math. Sem. Notes Kobe Univ.* **9** (1981), 75–84.
- [24] A. KAWAUCHI. *A Survey of Knot Theory* (Birkhäuser Verlag, 1996).
- [25] M. R. KELLY. A bound on the fixed-point index for surface mappings. *Ergodic Theory Dynam. Systems* **17** (1997), 1393–1408.
- [26] K. H. KIM, N. S. ORMES and F. W. ROUSH. The spectra of nonnegative integer matrices via formal power series. *J. Amer. Math. Soc.* **13** (2000), 773–806.
- [27] R. KIRBY. Problems in low-dimensional topology; *Geometric Topology* (W. H. Kazez, ed.), Studies in Advanced Mathematics (Amer. Math. Soc., 1997).
- [28] F. LAUDENBACH. Construction de difféomorphismes pseudo-Anosov. *Astérisque* **66–67** (1979), 243–250.
- [29] F. LAUDENBACH. La théorie de Thurston sur les surfaces à bord. *Astérisque* **66–67** (1979), 209–224.
- [30] D. H. LEHMER. Factorization of certain cyclotomic functions. *Annals of Math.* **34** (1933), 461–479.
- [31] C. J. LEININGER. On groups generated by two positive multi-twists: Teichmüller curves and Lehmer's number. *Geom. Topol.* **8** (2004), 1301–1359 (electronic).
- [32] R. C. LYNDON and P. E. SCHUPP. *Combinatorial Group Theory* (Springer-Verlag, 1977).
- [33] A. MANNING. Axiom A diffeomorphisms have rational zeta function. *Bull. London Math. Soc.* **3** (1971), 215–220.
- [34] T. MATSUOKA. Braids of periodic points and a 2-dimensional analogue of Sharkovskii's ordering. *Dynamical Systems and Nonlinear Oscillations* (Kyoto, 1985) 58–72. *World Sci. Adv. Ser. Dynam. Systems* **1** (World Scientific Publishing, 1986).
- [35] C. T. MCMULLEN. Polynomial invariants for fibered 3-manifolds and Teichmüller geodesics for foliations. *Ann. Sci. École Norm. Sup.* **33** (2000), 519–560.
- [36] H. MINC. *Nonnegative Matrices* (Wiley/Interscience, 1988).
- [37] M. J. MOSSINGHOFF. Polynomials with small Mahler measure. *Math. Comp.* **67** (1998), 1697–1705.
- [38] G. PÓLYA and G. SZEGŐ. *Problems and Theorems in Analysis II* (Springer-Verlag, 1976).
- [39] G. A. RAY. A locally parameterized version of Lehmer's problem. In *Mathematics of Computation 1943–1993: A Half-Century of Computational Mathematics* (W. Gautschi, ed.), *Proc. Symp. Appl. Math.* **48** (1994), 573–576.
- [40] D. ROLFSEN. *Knots and Links* (Publish or Perish, 1976).
- [41] E. RYKKEN. Expanding factors for pseudo-Anosov homeomorphisms. *Mich. Math. J.* **46** (1999), 281–296.
- [42] K. SCHMIDT. *Dynamical Systems of Algebraic Origin* (Birkhäuser Verlag, 1995).
- [43] D. S. SILVER and S. G. WILLIAMS. Mahler measure, links and homology growth. *Topology* **41** (2002), 979–991.
- [44] D. S. SILVER and S. G. WILLIAMS. Mahler measure of Alexander polynomials. *J. London Math. Soc.* **69** (2004), 767–782.
- [45] C. J. SMYTH. On the product of the conjugates outside the unit circle of an algebraic integer. *Bull. London Math. Soc.* **3** (1971), 169–175.
- [46] J. R. STALLINGS. Topologically unrealizable automorphisms of free groups. *Proc. Amer. Math. Soc.* **84** (1982), 21–24.
- [47] W. P. THURSTON. Three dimensional manifolds, Kleinian groups and hyperbolic geometry. *Bull. Amer. Math. Soc.* **6** (1982), 357–381.
- [48] W. P. THURSTON. *The Geometry and Topology of 3-Manifolds* (Lecture Notes, Princeton 1977. Revised version Princeton University Press, 1983).
- [49] W. P. THURSTON. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc.* **19** (1988), 417–431.
- [50] V. TURAEV. *Introduction to Combinatorial Torsions* (Birkhäuser Verlag, 2001).