


STRESS ULCER PROPHYLAXIS



Enteral famotidine or pantoprazole should be used whenever functioning GI is present

Dose adjust famotidine to daily for CrCl \leq 50 mL/min

Acute risk factors:

1. Mechanical ventilation without enteral nutrition
2. Chronic liver disease
3. Concerning coagulopathy (intrinsic INR > 1.5 or platelets < 50)
4. Burns > 35% TBSI
5. 2 or more of the following moderate risk factors:
 - a. Mechanical ventilation with enteral nutrition
 - b. AKI
 - c. Shock (MAP < 65)

References

ASHP Therapeutic Guidelines on Stress Ulcer Prophylaxis. ASHP Commission on Therapeutics and approved by the ASHP Board of Directors on November 14, 1998. *Am J Health Syst Pharm* 1999;56:347.

Guillamondegui, OD, Gunter OL et al. Practice management guidelines for stress ulcer prophylaxis, Eastern Association for the Surgery of Trauma (EAST), Chicago 2008. p. 24.

Marik PE, Vasu T, Hirani A, et al. Stress ulcer prophylaxis in the new millennium: a systematic review and meta-analysis. *Crit Care Med* 2010; 38:2222.

Stress ulcer prophylaxis. [SurgicalCriticalCare.net/AcuteCareSurgery.net](http://surgicalcriticalcare.net/AcuteCareSurgery.net). OrlandoHealth Surgical Critical Care and Acute Care Surgery Fellowships website. <http://surgicalcriticalcare.net/guidelines.php>. January 1, 2004. Updated September 15, 2011. Accessed July 23, 2015.

Ye Zhikang, Reintam Blaser Annika, Lytvyn Lyubov, Wang Ying, Guyatt Gordon H, Mikita J Stephen et al. Gastrointestinal bleeding prophylaxis for critically ill patients: a clinical practice guideline *BMJ* 2020; 368:l6722